OBSERVED

CLIMATE TRENDS

AND PROJECTED

CLIMATE CHANGE

IN THE PHILIPPINES





Republic of the Philippines

Department of Science and Technology (DOST)

Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA)

© PAGASA, 2018

ISBN 978-621-95882-0-1

Published by:

Philippine Atmospheric, Geophysical and Astronomical Services Administration Agham Rd., Diliman, Quezon City, 1100 Philippines

Tel.: +632-434-8130 Fax: +632-434-5882

Web site: www.pagasa.dost.gov.ph

Editors:

Flaviana D. Hilario, Edna L. Juanillo, and Rosalina G. De Guzman

Contributing Authors:

Thelma A. Cinco, Marcelino Q. Villafuerte II, Emma D. Ares, John A. Manalo, Wilmer A. Agustin, Kimberli Anne M. Aquino, and Ralph Q. Gaspar

Responsible Office at PAGASA:

Climatology and Agrometeorology Division (CAD)

Cover design and layout:

Wilmer A. Agustin

Referencing this report:

PAGASA, 2018: Observed and Projected Climate Change in the Philippines. Philippine Atmospheric, Geophysical and Astronomical Services Administration, Quezon City, Philippines, 36 pp.

Department of Science and Technology

Philippine Atmospheric, Geophysical and Astronomical Services Administration







Message



The Philippines is an archipelagic country situated in the northwestern region of the Pacific, making it highly exposed to natural hazards such as drought, earthquakes, floods, and tropical cyclones. Consequently, the country ranks third in the 2017 World Risk Index (WRI) of the United Nations University and fifth in the Long-Term Germanwatch Climate Risk Index (CRI) (1996-2015). It is therefore vital that we enhance our coping and adaptive capacities to natural hazards, particularly to climate-related threats that could be worsened by climate variability and change.

It has been seven years since the publication of the *Climate Change* in the *Philippines* document in February 2011, which has served as an important reference to various users of climate projections. With recent changes and innovations in the global field of climate study,

the Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA) has stepped up efforts to update its existing set of local climate information. Using the latest climate models, PAGASA has generated a collection of datasets from simulations of future climate conditions, termed as climate projections.

The Observed Climate Trends and Projected Climate Change in the Philippines – Policy Brief contains the basic findings of the past and present climate conditions and provides useful information from a range of plausible climate projections that are necessary to guide decision-makers in formulating strategic plans and policies for possible future climate conditions.

On behalf of PAGASA and our partners, I am honored to share the results of an extensive study on the Philippine climate.

VICENTE B. MALANO, Ph. D. Administrator, PAGASA



Preface

This report summarizes important findings about the state of the Philippine climate based on the most recent observations, as well as climate projections, which were derived from state-of-the-art climate models utilizing information from updated future climate scenarios. This report is very timely especially in the formulation of local climate change action plans and in the mainstreaming activities of the national and local government units. This may serve as a reference for different users needing information in formulating climate change adaptation strategies at the national and local levels.

For interested readers, a comprehensive guide in the proper utilization of the climate projections using PAGASA's newly developed Climate Information Risk Analysis Matrix (CLIRAM) is provided herein. Also included in this report are the summaries of the seasonal climate projection data for each province of the Philippines. We have high hopes that many users will find the information useful for their own applications, and we look forward to everybody's support and cooperation in promoting a climate-resilient country.

PAGASA extends its deepest gratitude to the following individuals, and the agencies they represent, for their expertise, tireless assistance, and motivation in the generation of the projection data from which this report was based upon: Joseph Daron, Richard Jones, Ian Macadam, Florian Gallo, Ron Kahana, Claire Scannell, David Hein, Simon Tucker, Rosanna Amato, and David Corbelli of the UK Met Office Hadley Center; Jack Katzfey and Dewi Kirono of Commonwealth Scientific and Industrial Research Organisation (CSIRO), Australia; Kevin Hodges of the Department of Meteorology, University of Reading, Reading, United Kingdom; and Sharon Taylor of the Philippine Rural Reconstruction Movement (PRRM).



Key Findings



Temperature

Observed temperature in the Philippines is warming at an average rate of 0.1°C/decade. Climate projections suggest continuous warming in the future. It is projected that the country-averaged mean temperature could increase by as much as 0.9°C–1.9°C (assuming the moderate emission scenario, RCP4.5) and 1.2°C–2.3°C (considering the high emission scenario, RCP8.5) in the mid-21st century (2036–2065). Warmer conditions are further expected by the end of the 21st century (2070–2099), which could range from 1.3°C–2.5°C (based on the RCP4.5) to 2.5°C–4.1°C (based on the RCP8.5) increase in mean temperature relative to the baseline climate.



Rainfall

Increasing trends in annual and seasonal rainfall were observed in many parts of the country. Such trends were found to be associated with extreme rainfall events. Multi-model projections suggest a range of increase and decrease in seasonal-mean rainfall exceeding 40% of its historical values. Nevertheless, the multi-model central estimate of projected changes in rainfall could be within the natural rainfall variations, except for the projected rainfall reduction over central sections of Mindanao that are beyond the observed rainfall variations in the past.



Tropical Cyclone

In the past 65 years (1951–2015), a slight decrease in the number of tropical cyclones (TCs) and a minimal increase in the frequency of very strong TCs (exceeding 170kph) were observed over the Philippine area of responsibility (PAR). These trends are projected to continue in the future. It has to be noted, however, that the high year-to-year variations in the frequency of occurrence and intensity of TCs remain to be dominant in the future Philippine climate conditions.



Sea Level Rise

The sea level has risen by nearly double the global average rate of sea level rise over certain parts of the Philippines from 1993 to 2015. Projections reveal that sea level in the country is expected to increase by approximately 20 cm by the end of the 21st century under the RCP8.5 scenario. Such projected increase in sea level might worsen storm surge hazards particularly on coastal communities.



Changes in Temperature

Observed temperature in the Philippines is warming at an average rate of 0.1°C/decade. Climate projections suggest continuous warming in the future. It is projected that the country-averaged mean temperature could increase by as much as 0.9°C–1.9°C (assuming the moderate emission scenario, RCP4.5) and 1.2°C–2.3°C (considering the high emission scenario, RCP8.5) in the mid-21st century (2036–2065). Warmer conditions are further expected by the end of the 21st century (2070–2099), which could range from 1.3°C–2.5°C (based on the RCP4.5) to 2.5°C–4.1°C (based on the RCP8.5) increase in mean temperature relative to the baseline climate.

Observed changes in temperature

Over the past 65 years (1951–2015), the annual mean temperature in the Philippines has risen by 0.68°C, which translates to an average rate of increase by about 0.1°C/decade (Fig. 1a). The country's annual maximum temperature, on the other hand,

has increased at a slower rate of approximately 0.05°C/decade while the annual minimum temperature has increased more rapidly at 0.15°C/decade (Fig. 1b and 1c, respectively).




Figure 1. Time series of observed (a) annual-mean, (b) annual-minimum, and (c) annual-maximum temperature anomalies in the Philippines. Anomaly values were taken relative to the 1971–2000 baseline period; thick black lines are the linear fit of the data indicating their long-term trends. All trends are found to be statistically significant at 5% level based on the Mann-Kendall trend test (see, Ref. 1).

Projected changes in temperature

The annual mean surface air temperature in the Philippines is expected to increase from 0.9°C to 1.9°C for the moderate emission scenario (RCP4.5) and from 1.2°C to 2.3°C for the high emission scenario (RCP 8.5) in the mid-21st century 2036–2065), and from 1.3°C to 2.5°C (RCP4.5) to as warm as 2.5°C to 4.1°C (RCP8.5) by the end of 21st century (2070–2099;




Figure 2. Time series of projected annual-mean temperature anomalies in the Philippines. Anomaly values were taken relative to the 1971–2000 baseline period. Shadings indicate the inter-model range. For comparison, the observed values from 1951 to 2015 are included (thick black line). The bars correspond to the 10th, 50th and 90th percentiles; while the whiskers correspond to the minimum and maximum projected temperature anomaly values by the mid- and end of the 21st century.





Projected changes in temperature


The spatial distribution of observed annual mean temperature is shown in Fig. 2a; most areas in the country have exprienced air temperatures exceeding 26°C, while, as expected, slightly cooler areas are found in

mountainous regions. These temperatures are projected to increase uniformly and minimally across the country in both the mid-21st century and the late 21st century (Fig. 2e-f).

Figure 3. Spatial distribution of (a) observed annual average temperature for 1971-2000 and projected temperature changes for the mid-21st century (2036-2065) at (b) RCP 4.5 and (c) RCP 8.5 relative to the aforementioned baseline period. Observed data is sourced from the Asian Precipitation - Highly Resolved Observational Data Integration Towards Evaluation (APHRODITE) project.







Changes in Rainfall

Increasing trends in annual and seasonal rainfall were observed in many parts of the country. Such trends were found to be associated with extreme rainfall events. Multi-model projections suggest a range of increase and decrease in seasonal-mean rainfall exceeding 40% of its historical values. Nevertheless, the multi-model central estimate of projected changes in rainfall could be within the natural rainfall variations, except for the projected rainfall reduction over central sections of Mindanao that are beyond the observed rainfall variations in the past.

Observed changes in rainfall

Changes in rainfall over the Philippines have been found to vary spatially^{2,3,4} and thus more effective to present in terms of spatial maps (Fig. 4). From 1951 to 2010, the annual total rainfall over northern sections of Luzon, Palawan, western sections of Visayas, and central and western sections of Mindanao have declined while increasing trends have

been observed in other areas, notably, the central parts of Luzon, eastern section of Visayas, and the northeastern and southwestern sections of Mindanao at a rate ranging from 10mm/decade to as much as a rate exceeding 40mm/decade. Such trends in annual total rainfall are associated with extreme rainfall events.




Figure 4. Observed trends in annual total rainfall and extreme rainfall in the Philippines during the period from 1951 to 2010. Extreme rainfall is defined here as the annual total rainfall taken from those days that exceeded the 95th percentile of rainfall on wet days (RR ≥1 mm) in the 1971–2000 baseline period. Areas marked with dots denote statistical significance at 5% level based on the Mann-Kendall test.



Rainfall in the Philippines is not only spatially variable, but also highly seasonal^{3,4}; hence, it is interesting to know whether different trends can be observed across seasons. Figure 5 shows the trends in seasonal total rainfall. Increasing trend in rainfall is observed pronouncedly over the northeastern section of Mindanao and eastern portions of Visayas in December-January-February coinciding to the northeast monsoon season. Note that the northeast monsoon season coincides also to the wet season over those areas implying for an increased risk of flooding potential on parts of the areas mentioned. Increasing trends in rainfall are also observed over central portions of Luzon and northeastern

sections of Mindanao in March–April–May season. In the following seasons of June–July–August and September–October–November, similar patterns of increasing rainfall trends are observed over the southern parts of Ilocos Region, and northeastern parts of Ilocos Region, and north eastern and southern parts of Mindanao; while decreasing trend is apparent in most parts of the country.

A noticeable drying trend can be observed over the northeastern portion of Luzon, and in central and northwestern sections of Mindanao in almost all seasons.




Figure 5. Observed trends in seasonal total rainfall in the Philippines during the period from 1951 to 2010. Areas marked with dots denote statistical significance at 5% level based on the Mann-Kendall test.



Projected changes in rainfall

The models used in this report are suggesting a wide range of future changes in Philippine rainfall (Fig. 6). Considering the models' assumptions underlying RCP8.5 scenario, the driest possible rainfall change (i.e., the 10th percentile of the models' projections) could reach beyond 40% reduction in many areas, particularly over Mindanao by the mid-21st century. The wettest possible change (i.e., the 90th percentile of the models' projections), on the other hand, could exceed a 40% increase in rainfall, particularly over

Luzon, western sections of Visayas, and some parts of Mindanao. The multi-model central estimate (i.e., the 50th percentile or the median) future rainfall conditions will be well within its natural variability (i.e., <1 standard deviation of observed rainfall); except for the drier future over central sections of Mindanao, particularly in September-October-November and the December-January-February seasons, which might require actionable climate change adaptation plans.




Figure 6. Projected change in seasonal total rainfall in the Philippines for the mid-21st century (2036–2065) relative to the baseline period of 1971–2000. The wettest possible change represents the 90th percentile of the projections, 50th percentile is the median or usual change, and 10th percentile is driest possible change; these were computed from the 12-member high emission scenario (RCP8.5) RCM ensemble. Dotted areas denote that the change beyond natural variability (i.e. ±1 standard deviation of the baseline period).



Changes in Tropical Cyclone

In the past 65 years (1951–2015), a slight decrease in the number of tropical cyclones (TCs) and a minimal increase in the frequency of very strong TCs (exceeding 170kph) were observed over the Philippine area of responsibility (PAR). These trends are projected to continue in the future. It has to be noted, however, that the high year-to-year variations in the frequency of occurrence and intensity of TCs remain to be dominant in the future Philippine climate conditions.

Observed changes in tropical cyclones

The number of tropical cyclones (TCs) entering the Philippine area of responsibility (PAR), and the number of TCs that made landfall, shows a minimal decreasing trend from 1951 to 2015 (Fig. 7a). Looking at the number of very strong TCs with

maximum sustained wind speeds exceeding 170 kph, a slight increasing trend is observed during the period from 1980 to 2015 (when more reliable and consistent observations were taken), as shown by the red dashed line in Fig. 7b.




Figure 7. (a) Time series of annual number of TCs over PAR and landfalling TCs from 1951 to 2015, (b) time series of annual count of all (black line) and very intense tropical cyclones (maximum sustained winds >170kph, red line) that entered the PAR from 1980 to 2015, and (c) kernel density plots estimated from maximum sustained wind speeds of all the tropical cyclones that existed in PAR comparing two time periods (1980-1997 and 1998-2015, black and red curves, respectively). Dashed lines in (a) and (b) indicate linear trends. The TC data shown in (a) were taken from PAGASA while (b) and (c) were based from the Japan Meteorological Agency.



Projected changes in tropical cyclones

Scientists from PAGASA and the UK Met Office has worked together to investigate the possible changes in tropical cyclones that could affect the Philippines in the future⁵. Considering the five regional climate model simulations, which cover a domain that includes the PAR and assuming a high emission scenario (RCP8.5), the projected changes in the behaviour of TCs in the mid-21st century (2036-2065) were found consistent with the currently observed trends (Fig. 8). Three of the models suggest that decrease in tropical cyclone frequency is significant, while the two others suggest that no change is expected. In terms of the TC intensity, four of the models agree in a projected increase, two of which are significant. This is consistent with the report of the IPCC that in the 21st century the average annual number of TCs in the Western North Pacific is expected to decrease, and that an increase in the frequency of strong TCs in the region is "more likely than not".

A caveat in the projections is that not all possible contributing factors in the development and behaviour of TCs were considered. The model simulations further indicate that the year-to-year variability will remain high in the future⁵. Hence, the lessons we learned in the past in mitigating tropical cyclone impacts are important to consider in the formulation of plans and programs for climate change adaptation.

		Climate	Model Sim	ulations			
	1	2	3	4	5		
Change in tropical cyclone frequency	+	1	-	-	+		
Change in tropical cyclone intensity	-	1	1	1	*		

Figure 8. Projected changes in tropical cyclones affecting the Philippines by the mid-21st century assuming high emission scenario (RCP8.5). Black arrows indicate significant changes, grey arrows indicate insignificant changes, and a dash indicates no change. Taken from [5].



Changes in Sea Level

The sea level has risen by nearly double the global average rate of sea level rise over certain parts of the Philippines from 1993–2015. Projections reveal that sea level in the country is expected to increase by approximately 20 cm by the end of the 21st century under the RCP8.5 scenario. Such projected ease in sea level might worsen storm surge hazards particularly on coastal

communities.

Observed changes in sea level

Based on satellite observations (AVISO altimetry data) taken from 1993 to 2015, the sea level has risen by as much as 5.7–7.0 mm/yr over the Philippine Sea (Fig. 9). Such rate is approximately double the highest global average rate of 2.8–3.6 mm/yr, which was observed between 1993 to 2010. This difference could be attributed to the occurence of natural climate-related phenomena, e.g. El Niño Southern Oscillation (ENSO), which directly affects the tropical Pacific region⁷.

Looking closely at different coastal areas in the country, a rate of sea level rise as high as 4.5–5.0 mm/yr is observed east of the islands of Leyte and Samar, and along the south western coasts of the Central and Western Visayas, and east of Mindanao and south of Zamboanga (Fig. 9).

Noticeable changes were also observed at the local level (Fig. 10). Tide gauge observations from National Mapping and Resource Information Authority (NAMRIA) indicate that a rapid increase in sea level was observed in Manila, particularly from 1955 to 2015. However, this is attributed to long-term land subsidence from excessive groundwater




Figure 9. Sea level changes in the Philippines region from 1993-2015. Taken from [9].

extraction⁸. Gradual increases in sea level were observed in Legazpi and Davao, while no apparent trend was observed in Cebu and Jolo, Sulu.




Figure 10. Sea level rise in selected areas (in mm) above the Revised Local Reference (RLR) level (7,000 mm). Data source: National Mapping and Resource Information Authority (NAMRIA)

Projected changes in sea level

As shown in Figure 11, sea level rise in the Philippines will continue to be slightly larger than the global average⁹. In both the moderate (RCP4.5) and high (RCP8.5) emission scenarios, the increase is expected to be almost the same by the mid-21st century. The trend for RCP4.5 will continue to be linear up to the end of the 21st century, while the trend for RCP8.5 will follow a

rather exponential increase - leading to a sea level rise by approximately 20 cm (Fig. 11b and 11c). The projected increase in sea level might worsen storm surge hazards particularly on coastal communities, which requires climate change adaptation action plans.




Figure 11. (a) shows the historical sea level change (SLC) from tide gauge observations and future projections; (b) and (c) contain time series plots of the global mean sea level change projections under RCP4.5 and RCP8.5. The plot in (a) shows the sum of SLC components for the global ocean (black line, with uncertainty range in grey) and for the coastal region of Legaspi City, Philippines (blue line, with uncertainty), under the RCP8.5 future scenario. Annual mean tide-gauge measurements from Legaspi (blue line) and Cebu (green line) are also presented and a linear trend is added to the observations from Legaspi. The solid line in (a), (b), and (c) represent the central estimate and the shaded area represent the uncertainty of the likely range. Taken from [9]



Methodology

This section contains a brief discussion on how climate projections for the Philippines were generated — the models utilized, the model settings, the time slices, and the range of possible climate futures. The latest climate scenarios, the Representative Concentration Pathways (RCPs), are also introduced.

The previous climate change information for the Philippines were presented in the Climate Change in the Philippines by PAGASA on 2011¹⁰. These were based on future socio-economic conditions known as the Special Report Emission Scenario (SRES). The SRES covered a wide range of possible future atmospheric GHG concentration, however none were based on policies for climate change mitigation. To resolve this, the IPCC released a new set of scenarios representing the cumulative concentration of GHGs. These new set of scenarios were designated as the

Representative Concentration Pathway
Special Report Emission
Scenarios

simulates atmospheric-oceanic processes

100-km resolution

25-km resolution

Figure 12. Illustration of the general scheme of the climate modeling processes that produced the currently available climate projections for the Philippines [9].

Regional Climate Model

Representative Concentration Pathways (RCPs) and have names based on radiative forcing (energy accumulated by the GHGs in the atmosphere) levels. The main difference between the new RCPs and the previous SRES scenarios is that there are no fixed sets of assumptions related to population growth, economic development, or technology associated with any RCP. Many different socio-economic futures are possible leading to the same level of radiative forcing.

The new set of climate projections for the Philippines were based on two of the most recent scenarios from the IPCC: RCP4.5 (moderate level of GHG emissions) and RCP8.5 (high level of GHG emissions).

Figure 12 illustrates how the climate projections were derived. Initially, the global climate models (GCMs) project the possible future climate conditions given a particular greenhouse gases concentration (i.e., the RCPs). Different physical and dynamical processes of the climate system are represented in the GCMs. The GCM-derived climate projections are then downscaled by the regional climate models (RCMs), namely the Conformal Cubic Atmospheric Model (CCAM)¹¹, the Providing Regional Climates for Impacts Studies (PRECIS) model¹², the RegCM4¹³, and the HadGEM3-RA¹⁴, to provide more realistic and spatially detailed quantities (e.g., temperature, pressure, etc.) at local scales (i.e. downscaled)¹⁵. For а technically detailed discussion of all the models used in generating the projections, refer to [16].





The CLIRAM Tool

This section introduces the Climate Information Risk Analysis Matrix tool and provides a comprehensive guide on its utilization

Using the collection of RCP-based projections available for the Philippines, PAGASA has derived a range of climate futures to help policy makers, such as LGUs, establish robust decisions on mitigation and adaptation options¹⁷. This new approach in utilizing climate projection data is the underlying concept in the Climate Information Risk Analysis Matrix (CLIRAM) tool.

The CLIRAM provides the projected changes in climate variables (particularly for rainfall, mean temperature, minimum temperature, and maximum temperature) in both the mid-21st century (2036-2065) or the late-21st century (2070-2099) based on the 1971–2000 baseline period. Following the desciptions of the parts of the CLIRAM and an instruction on its utilization which was partly adapted from the procedures found in the Climate and Disaster Risk Assessment (CDRA) process.

Parts of the CLIRAM Tool

The current version of the CLIRAM tool is based on the suggestions and needs of participants in the pilot testings. Originally, there were separate CLIRAMs for each season. The new layout (see Tables 1 and 2) displays data for all seasons, which is helpful for the analysis and interpretation of interseasonal patterns of change. Column 1 in both Tables indicate the season as well as the observed baseline value for the period (i.e. from 1971 to 2000). Projections for each season are divided into two sets based on the the driving scenario of the climate model sources (see Column 2). These projections are further clustered into percentiles, categorized in Column 3 as the lower bound (10th percentile), median (50th

percentile), and upper bound (90th percentile). Column 4 cointains the amount of change between the target period and the baseline period in terms of algebraic difference in the temperature CLIRAMs and percent difference in the rainfall CLIRAM (which is provided as actual change amount in Column 5). The projected values for the target period is listed on Column 5 and Column 6 of the temperature (i.e. for the minimum, maximum, and mean) and rainfalll CLIRAM, respectively. Instructions on filling up the blank columns are provided are as follows.

Utilizing the CLIRAM Tool

In accomplishing the CLIRAM for seasonal total rainfall, determine the projected patterns of change (Column 6 of Table 1) based on the following (refer to Column 4 to 6):

- 1. The lower bound that indicates the driest possible change;
- 2. The upper bound that indicates the wettest possible change; and
- 3. The median which represents the most likely future change.

If the reader is familiar with CDRA, one may notice in Table 1 that unlike Task 1 of Step 1 in the CDRA process, the CLIRAM does not require any computation, but instead already provides the values necessary for Columns 5 and 6. The same goes for the mean temperature CLIRAM (see Table 2).



			Project	Projected Change	Projected			
Season	Scenario	Range*	Percent (%)	Rainfall amount (mm)	Seasonal Rainfall Amount (mm)	Information about patterns of change**	Potential Impacts	Adaptation Option
Col 1	Col 2	Col 3	Col 4	Col 5	Col 6	Col 7	Col 8	Col 9
		Lower Bound	-0.1	-0.2	107.3	Minimal to no change		
	Moderate Emission	Median	17.71	19.0	126.5	Rainfall will most likely increase (17 %) during the Northeast (NE) Monsoon or Amihan As	Assess the	
December-January- February (DJF)	(RCP4.5)	Upper Bound	55.5	59.6	167.1	The wettest possible rainfall change during the Northeast (NE) Monsoon or Amihan USI shows an increase of 55%. This inrease could be beneficial to some sectors of the Site community, as this season corresponds to the dry months over the region CC	potential impacts using the Steps 2-4 of the CDRA process	Assess the
		Lower Bound	2.7	2.9	110.4	Minimal to no change		potential impacts
Observed baseline = 107.5 mm	High Emission	Median	27.8	29.9	137.4	Rainfall will most likely increase (20%)during the Northeast (NE) Monsoon or Amihan, same with the RCP4.5		Steps 5-6 of the CDRA process
	(RCP8.5)	Upper Bound	53.4	57.4	164.9	The wettest possible rainfall change during the Northeast (NE) Monsoon or Amihan shows an increase of 53%. This inrease could be beneficial to some sectors of the community, as this season corresponds to the dry months over the region		
		Lower Bound	0.7	1.3	199.8	Minimal to no change		
	Moderate Emission	Median	6.9	13.7	212.2	Minimal to no change		
March-April-May	(NCP4.3)	Upper Bound	25.7	51.1	249.6	Increase in rainfall during the dry months		
(MAM) Observed baseline =	High Emission	Lower Bound	-7.7	-14.3	184.2	A driest possible change in rainfall shows a slight reduction of about 7% during the dry season. This change could have adverse impacts on human health, water resources, and businesses, among others.		
	(NCF0.3)	Median	4.8	9.6	208.1	Minimal to no change		
		Upper Bound	19.8	39.4	237.9	Increase of 20% in rainfall during the dry season,		
	Moderate Emission (RCP4.5)	Lower Bound	-21.3	-249.4	920.8	The driest possible rainfall change during the Southwest Monsoon (SW) or Habagat shows a 20% reduction. Areas already vulnerable to flooding will continue to be affected, but this might affect the availability of water resources.		
June-July-August		Median	-10.1	-118.6	1051.6	Minimal to no change		
		Upper Bound	-0.4	-5.0	1165.2	Minimal to no change		
Observed baseline = 1170.2 mm	High Emission (RCP8.5)	Lower Bound	-17.0	-199.3	970.9	The driest possible future rainfall change during the Southwest Monsoon (SW) or Habagat shows a 17% reduction. Areas already vulnerable to flooding will continue to be affected, but this might affect the availability of water resources.		
		Median	-6.1	-71.7	1098.5	Minimal to no change		
		Upper Bound	/:/	90.3	1260.5	Minimal to no change		
	Moderate Emission	Lower Bound	-10.8	-82.3	676.4	The driest possible future rainfall change during the transition period from NE to SW monsoon shows a reduction of 10%.		
September-October- November (SON)	(RCP 4.3)	Median	-6.0	-45.2	713.5	Minimal to no change		
(100)		Upper Bound	7.7	58.5	817.2	Minimal to no change		
Observed baseline =		Lower Bound	-8.0	9.09-	698.1	Minimal to no change		
758.7 mm	High Emission	Median	3.9	29.6	788.3	Minimal to no change		
	(RCP 8.5)	Upper Bound	19.9	150.7	909.4	The wettest possible rainfall change is about 19%, during the transition period		

Table 1. Sample CLIRAM of projected changes in seasonal total rainfall in the mid-21st century (2036-2065) for Metro Manila; baseline period = 1971-2000.

bound: 90th percentile; median: 50th percentile; lower bound: 10th percentile



For each season, assess first the lower bound that represents the lowest possible rainfall change. Determine whether the reduction is relevant relative to the baseline climate. For example in Table 1, on the March–April–May season, the lower bound change (with red highlight) under the high emission scenario shows a slight rainfall reduction. Considering that this projection is on the dry season of Metro Manila, it could have potential impacts to various sectors.

The same procedure can be used to assess the seasonal mean temperature in Table 2. Identify the lower and upper bound to get the temperature range for each season and input this on Column 6. Based on these, identify the coldest and hottest seasons. The projected temperature shows warming in all seasons, but a greater warming is indicated under the high emission scenario (RCP8.5).

For users to assess the potential impacts (see Column 8 of Table 1 and Column 7 of Table 2) on their respective sectors for each of the projected change, they could refer to steps 2–4 in the CDRA process. In determining the impacts, a particular climate risk must be identified. For example, if one is looking at risk of flooding, then consider the wettest possible change during the wet season.

Possible adaptation options for each potential impact should be identified following Steps 5 & 6 in the CDRA process. Ideally, this part should be carried out by the focal lead from the LGU (either the Planning Officer, Local Disaster Risk Reduction Officer, or Municipal Agriculturist).

Examples of accomplished CLIRAMs of both seasonal rainfall and mean temperature in the mid-21st century are shown in the Appendix. These were filled-out by officials from the Municipality of Salcedo, Eastern Samar during the conduct of a CLIRAM utilization workshop.

A summary of the seasonal climate projections for all Philippine provinces are provided in the Annex. To aid in the analysis, a provincial-level map of the distribution of the projected changes is also available. If interested to acquire the detailed CLIRAM of a particular area/s (e.g. the example provided herein) including the projection maps, the reader may request from PAGASA through any of the contact information provided at the back portion of the cover page of this report.



			Projecte	Projected Change	Information		
Season	Scenario	Range*	Degrees Celsius	Degrees Celsius	about patterns of change	Potential Impacts	Adaptation Option
Col 1	Col 2	Col 3	Col 4	Col 5	9 loo	Col 7	Col 8
		Lower Bound	1.0	27.1	Coldest season		
December-January-	(RCP 4 5)	Median	1.2	27.3	27.1 to 27.7		
February (DJF)	(Cit Iswi)	Upper Bound	1.6	7.72		Assess the	
Observed baseline =		Lower Bound	1.2	27.3		using the	
26.1 °C	(PCP 8 5)	Median	1.6	7.72	27.3 to 28.0	Steps 2-4 of the	
	(C.6 TON)	Upper Bound	1.9	28.0		CDKA process	Od+ 00000 A
	Action of the second	Lower Bound	6.0	29.7			potential impacts
March-April-May	(RCP 4 5)	Median	1.2	30.0	29.7 to 30.5		using the
(MAM)	(C:t ISW)	Upper Bound	1.7	30.5			Steps 5-6 of the CDRA process
Children bearing		Lower Bound	1.3	30.1	Hottest season		
28.8°C	High Emission	Median	1.6	30.4	30.1 to 31.0		
	(co pu)	Upper Bound	2.2	31.0			
		Lower Bound	1.0	29.0			
June-July-August	Moderate Emission	Median	1.3	29.3	29.0 to 29.8		
(JJA)	(cut land	Upper Bound	1.8	29.8			
- orilozed bornoodO		Lower Bound	1.3	29.3			
28.0°C	High Emission (RCP & 5)	Median	1.5	29.5	29.3 to 30.3		
		Upper Bound	2.3	30.3			
	Modernto Emirrion	Lower Bound	1.0	28.4			
September-October-	(RCP 4.5)	Median	1.1	28.5	28.4 to 29.2		
November (SON)		Upper Bound	1.8	29.2			
Observed baseline =	1 Calculation	Lower Bound	1.3	28.7			
27.4°C	(RCP 8 5)	Median	1.5	28.9	28.7 to 29.0		
		Upper Bound	2.2	29.6			

Table 2. Sample CLIRAM of projected changes in seasonal mean temperature in the mid-21st century (2036-2065) for Metro Manila; baseline period = 1971-2000.

* upper: 90th percentile; median: 50th percentile; lower: 10th percentile



References

- 1. H.B. Mann. Nonparametric tests against trend. Econometrica 13, 245–259. (1945).
- 2. T.A. Cinco, R.G. de Guzman, F.D. Hilario, and D.M. Wilson. Long-term trends and extremes in observed daily precipitation and near surface air temperature in the Philippines for the period 1951–2010. Atmospheric Research 145–146, 12–26. (2014).
- M.Q. Villafuerte II, J. Matsumoto, I. Akasaka, H.G. Takahashi, H. Kubota, and T.A. Cinco. Long-term trends and variability of rainfall extremes in the Philippines. Atmospheric Research 137, 1–13. (2014).
- M.Q. Villafuerte II, J. Matsumoto, and H. Kubota. Changes in extreme rainfall in the Philippines (1911–2010) linked to global mean temperature and ENSO. International Journal of Climatology 35, 2033–2044.
- J. Daron, I. Macadam, F. Gallo, E. Buonomo, M. Villafuerte, T. Cinco and R. Jones. Future Tropical Cyclone Risk in the Philippines -Information Brief. Met Office, Exeter, Devon, United Kingdom. 1-38. (2016).
- IPCC. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 1535. (2013).
- B. Meyssignac, D. Salas Y Melia, M. Becker, W. Llovel, and A. Cazenave. Tropical Pacific spatial trend patterns in observed sea level: Internal variability and/or anthropogenic signature? Climate of the Past 8, 787–802. (2012).

- 8. K. S. Rodolfo and F. P. Siringan. Global sealevel rise is recognised, but flooding from anthropogenic land subsidence is ignored around northern Manila Bay, Philippines. Disasters 30, 118–139. (2006).
- R. Kahana, R. Abdon, J. Daron, C. Scannell, and M. Palmer. Projections of mean sea level change for the Philippines. Met Office, Exeter, Devon, United Kingdom and Phillippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA), Quezon City, Philippines. 29. (2016).
- PAGASA. Climate Change in the Philippines. Phillippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA), Quezon City, Philippines. 85. (2011).
- J. J. Katzfey, J. L. McGregor, and R. Suppiah. High-Resolution Climate Projections for Vietnam: Technical Report. Commonwealth Scientific and Industrial Research Organisation (CSIRO), Australia. 226. (2014).
- R. G. Jones, M. Noguer, D. C. Hassell, D. Hudson, S. S. Wilson, G. J. Jenkins, and J. F. B. Mitchell. Generating high resolution climate change scenarios using Precis. National Communications Support Unit Handbook. 40. (2004).
- F. Giorgi and co-authors. RegCM4: Model description and preliminary tests over multiple CORDEX domains. Climate Research 52, 7–29. (2012).
- G. M. Martin and co-authors. The HadGEM2 family of Met Office Unified Model climate configurations. Geoscientific Model Development 4, 723–757. (2011).



- 15. HLURB. CLUP Guidebook: A Guide to Comprehensive Land Use Plan Preparation. Supplemental Guidelines on Mainstreaming Climate and Disaster Risks in the Comprehensive Land Use Plan. Housing and Land Use Regulatory Board (HLURB), Quezon City, Philippines. 243. (2015).
- M. Villafuerte II and co-authors. Dynamical downscaled multi-model projections of future changes in rainfall and temperature over the Philippines. Submitted to Theoretical and Applied Climatology. (2017).
- J. Daron and co-authors. Providing future climate projections using multiple models and methods: insights from the Philippines. Submitted to Climatic Change. (2017).



Appendix A

Provided here are samples of accomplished CLIRAMs of seasonal total rainfall and seasonal mean temperature of the mid-21st century for the Province of Eastern Samar. Accompanying each CLIRAM are spatial maps of the distribution of rainfall and temperature in the province according to the 90th percentile, 50th percentile, and 10th percentile of the range of projection values.

			Projected Change	Change	Projected			
Season	Scenario	Range*	Percent (%)	Rainfall amount (mm)	Seasonal Rainfall Amount (mm)	Information about patterns of change	Potential Impacts	Adaptation Option
Col 1	Col 2	Col 3	Col 4	Col 5	Col 6	Col 7	Col 8	6 loo
		Lower Bound	-4.2	-41.1	945.9	Minimal to no change		
		Median	1.3	12.5	999.5	Minimal to no change		
December-January February (DJF)	Moderate Emission (RCP 4.5)	Upper Bound	45.0	444.4	1431.4	The highest possible future rainfall change during the Northeast (NE) Monsoon or Amihan shows an increase of 45%. This increase could be detrimental to some sectors of the community, as this season corresponds to the wettest months over the region	Farm flooding, crops ruined (e.g. wash out planted cassava 6-8 f months growing period) Pest & disease infestation Increased cost of lodging animals & crops Decrease in yield/production	
		Lower Bound	-7.9	-77.8	909.2	Minimal to no change		
Observed baseline =		Median	13.7	135.2	1122.2	Minimal to no change		
987 mm	High Emission (RCP 8.5)	Upper Bound	43.6	430.8	1417.8	The highest possible future rainfall change during the Northeast (NE) Monsoon or Amihan shows an increase of 44%. This increase could be detrimental to some sectors of the community, as this season corresponds to the wettest months over the region	Farm flooding, crops ruined (e.g. wash out planted cassava 6-8 months growing period) Pest & disease infestation Increased cost of lodging animals & crops Decrease in yield/production	
	Moderate	Lower Bound	-2.6	-12.1	452.0	Minimal to no change		
March-April-May	Emission	Median	1.8	8.2	472.3	Minimal to no change		
(MAM)	(RCP 4.5)	Upper Bound	17.4	80.8	544.9	Minimal to no change		
Observed baseline =	High Emission	Lower Bound	5.8	26.8	490.9	Minimal to no change		
464 mm	(PCD 9 5)	Median	1.1	5.3	469.4	Minimal to no change		
	(nor 0.3)	Upper Bound	13.0	60.3	524.4	Minimal to no change		
	Moderate	Lower Bound	-6.7	-37.5	522.3	Minimal to no change		
June-July-August	Emission	Median	-1.1	-6.0	553.8	Minimal to no change		
(NIA)	(RCP 4.5)	Upper Bound	8.3	46.4	606.2	Minimal to no change		
Observed baseline =	High Emission	Lower Bound	-14.0	-78.3	481.5	Minimal to no change		
260 mm	(RCP 8.5)	Median	-0.6	-3.2	556.6	Minimal to no change		
		Opper Bound	9.7	54.6	614.4	Minimal to no change		
	Moderate	Lower Bound	-20.1	-174.9	696.5	The driest possible future rainfall change during the transition period from NE to SW monsoon shows a reduction of 20%.		
September-October- November (SON)	(RCP 4.5)	Median	-16.6	-144.7	726.7	The most likely future rainfall change during the transition period from NE to SW monsoon shows a reduction of 17%		
		Upper Bound	-3.8	-33.4	838.0	Minimal to no change		
Observed baseline = 871 mm	High Emission	Lower Bound	-22.0	-191.3	680.1	The driest possible future rainfall change during the transition period from NE to SW monsoon shows a reduction of 22%		
	ווים שיוו	Median	-9.3	-81.3	790.1	Minimal to no change		
		Upper Bound	4.7	40.7	912.1	Minimal to no change		

(2036–2065); baseline period: 1971–2000.

Table A-1. Accomplished CLIRAM of the projected seasonal total rainfall in Eastern Samar for the mid-21st century






Figure A-1. Projected change in seasonal total rainfall in the Eastern Samar for the mid-21st century (2036–2065) relative to the baseline period of 1971–2000. The wettest possible change represents the 90th percentile of the projections, 50th percentile is the median or usual change, and 10th percentile is driest possible change; these were computed from the 12-member high emission scenario (RCP8.5) RCM ensemble.



Table A-2. Accomplished CLIRAM of the projected seasonal mean temperature for Eastern Samar in the mid-21st century (2036–2065); baseline period: 19712000.

			Proje	Projected Change	Information		
30000	Circus Co.	*		Projected Seasonal	about	Potential	Adaptation
Dospac	Scenario	Range	Change in °C	Mean Temperature	patterns of	Impacts	Option
				(₀ ,	change		
Col 1	Col 2	Col 3	Col 4	Col 5	Col 6	Col 7	Col 8
	Moderate	Lower Bound	1.0	27.1	Coldest season	 Impotent livestock/crops 	Conduct Climate Field
December-January-	Emission	Median	1.1	27.2	27.1 to 27.7	Investock disease	
February (DJF)	(RCP 4.5)	Upper Bound	1.6	7.72		Ice-ice effect	more resilient to climate
	a ciccian de il	Lower Bound	1.3	27.4		• Fish kill	adversities
Observed baseline =	HIGH EMISSION	Median	1.5	27.6	27.4 to 28.0	Doiv ai ageaga	Browide EWE for
26.1 °€	(KCP 8.5)	Upper Bound	1.9	28.0		production	agriculture
	Moderate	Lower Bound	1.0	28.7			• Drawido incursoso to
March-April-May	Emission	Median	1.2	28.9	28.7 to 29.4		formati
(MAM)	(RCP 4.5)	Upper Bound	1.7	29.4			מו וותו א
	I limb Caning	Lower Bound	1.4	29.1			
Observed baseline =	Ingh Emission	Median	1.6	29.3	29.1 to 29.8		
م-1.1 /2	(LC 0.3)	Upper Bound	2.1	29.8			
	Moderate	Lower Bound	1.0	29.3			
June-July-August (JJA)	Emission	Median	1.2	29.5	29.3 to 30.1		
	(RCP 4.5)	Upper Bound	1.8	30.1			
Observed baseline =	Ligh Emission	Lower Bound	1.4	29.7	Hottest season		
28.3°C	(PCB 8 E)	Median	1.7	30.0	29.7 to 30.5		
	(RCP 0.3)	Upper Bound	2.2	30.5			
	Moderate	Lower Bound	1.0	28.7			
September-October-	Emission	Median	1.2	28.9	28.7 to 29.5		
November (SON)	(RCP 4.5)	Upper Bound	1.8	29.5			
	Link Emission	Lower Bound	1.4	29.1			
Observed baseline =	Ingii Lillission	Median	1.5	29.2	29.1 to 29.9		
2/17	(RCP 8.5)	Upper Bound	2.2	29.9			

* upper: 90th percentile; median: 50th percentile; lower: 10th percentile






Figure A-2. Projected change in seasonal mean temperature in the Eastern Samar for the mid-21st century (2036–2065) relative to the baseline period of 1971-2000. The upper bound of the range of change represents the 90th percentile of the projections, 50th percentile is the median or usual change, and 10th percentile is lower bound of the range of change; these were computed from the 12-member high emission scenario (RCP8.5) RCM ensemble.



Appendix B

This section contains the summary of the seasonal values of total rainfall and mean temperature taken from the CLIRAM of each of the provinces in the Philippines. Note that currently there are no available climate projections for provinces of relatively small land areas since the resolution of most of the models are limited to $25 \times 25 \text{ km}$.

Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000.

		Obs	erved (1971-20	00)			-	Pro	jected (2	036-2065)				
9	3				-			DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Province		DJF	МАМ	JJA	SON	Scenario	Range*	Percent change	Projected value						
						Moderate	Lower Bound	1.5	50.5	-15.6	156.5	-23.9	841.5	-18.9	483.1
te	9					Emission	Median	12.8	56.2	-2.7	180.4	-19.9	886.5	0.4	597.7
No	2	40.0	105.5	1 100 1	595.4	(RCP4.5)	Upper Bound	32.2	65.8	22.4	227.1	10,4	1,221.0	29.8	772.7
llocos Norte	3	49.8	185.5	1,106.4	393.4	High	Lower Bound	-2.8	48.4	-25.9	137.5	-15.9	930.9	-14.9	506.6
100	3					Emission	Median	9.6	54.6	-8.1	170.5	-10.1	995.1	-2.2	582.5
						(RCP8.5)	Upper Bound	24.5	62.0	17.1	217.2	17.3	1,298.2	29.5	770.9
						Moderate	Lower Bound	-11.8	15.4	-10.3	259.2	-14.3	1,350.5	-19.1	544.3
Sur	5					Emission	Median	-2.7	17.0	-4.9	274.6	3.5	1,630.5	-5.2	637.6
SS	2	17.5	288.8	1,575.4	672.9	(RCP4.5)	Upper Bound	51.6	26.5	18.6	342.4	7.9	1,699.2	14.8	772.6
Ilocos		-		- Action (High	Lower Bound	-23.6	13.4	-21.7	226.0	-12.5	1,378.6	-11.3	596.9
	Ì					(RCP8.5)	Median	-0.2	17.5	-3.2	279.7	1.3	1,596.5	0.0	672.9
	-					(KCF0.3)	Upper Bound	25.4	21.9	25.6	362.7	19.1	1,876.8	12.7	758.5
						Moderate	Lower Bound	-0.1	14.7	-10.3	354.7	-13.9	1,594.9	-15.7	706.3
100						(RCP4.5)	Median	9.1	16.0	-2.6	385.3	5.2	1,949.1	4.7	877.5
La Union		14.7	395.6	1,852.3	837.8	(NCC 413)	Upper Bound	45.4	21.4	16.4	460.5	9.3	2,025.0	13.3	949.5
-53	3	1100		77.00		High	Lower Bound	-22.6	11.4	-20.4	314.8	-8.8	1,689.2	-17.3	693.2
In	П					(RCP8.5)	Median	14.8	16.9	-2.9	384.2	2.0	1,889.9	3.7	868.4
-	+		_			(00-00-)	Upper Bound	46.3	21.5	24.8	493.6	25.8	2,329.4	24.6	1,043.9
						Moderate	Lower Bound	4.8	20.3	-3.4	287.8	-22.3	1,249.5	-16.3	592.8
Pangasinan						(RCP4.5)	Median	10.8	21.5	-1.3	294.2	-13.3	1,395.0	-0.6	703.3
385	5	19.4	298.0	1,608.9	707.8	Comment	Upper Bound	41.6	27.5	17.7 -9.7	350.7	1.6	1,634.3	13.5 -3.7	803.5
Pane	5			1		High	Lower Bound	-8.9 9.6	17.7 21.3	0.6	269.2 299.8	-16.8 -3.3	1,338.4	0.9	681.7 713.9
F						(RCP8.5)	Median	55.7	30.2	27.5	379.9	9.9	1,556.4	17.7	832.8
	+					Or or other	Upper Bound Lower Bound	3.9	45.2	-16.3	184.7	-28.6	870.4	-23.4	485.7
	ш					Moderate Emission	Median	7.7	46.9	-4.4	210.8	-14.5	1,042.7	-4.3	607.3
	,			4.00		(RCP4.5)	Upper Bound	36.2	59.2	14.8	253.3	3.9	1,266.3	13.3	719.1
Abra		43.5	220.6	1,218.9	634.4	I	Lower Bound	2.0	44.4	-23.4	168.9	-23.5	932.7	-14.0	545.9
1						High Emission	Median	10.3	48.0	-8.2	202.6	-9.4	1,104.8	-5.8	597.9
						(RCP8.5)	Upper Bound	19.2	51.9	16.8	257.7	14.7	1,397.7	18.3	750.4
	1					45.14.1.23	Lower Bound	1.7	147.1	-7.2	170.8	-33.1	550.2	-12.3	631.3
	ш					Moderate Emission	Median	7.6	155.5	1.8	187.3	-21.6	645.4	5.5	759.9
08/	3		0200	0.000	1224	(RCP4.5)	Upper Bound	38.4	200.1	25.7	231.2	8.7	894.3	15.8	833.9
Apavao		144.6	184.0	822.7	720.1	Ulah	Lower Bound	-2.2	141.4	-14.8	156.7	-31.2	565.8	-11.4	638.2
	1					High Emission	Median	9.3	158.1	1.7	187.0	-1.9	807.4	4.3	750.9
						(RCP8.5)	Upper Bound	27.4	184.2	26.8	233.2	15.6	951.0	15.6	832.7
						Moderate	Lower Bound	6.5	50.8	-11.0	376.0	-24.6	1,308.3	-24.8	700.5
	Ш					Emission	Median	9.7	52.3	-3.3	408.4	-15.4	1,467.9	-6.1	875.0
one		47.7	422.2	1 724 0	931.8	(RCP4.5)	Upper Bound	35.6	64.7	27.0	536.2	5.9	1,836.6	15.6	1,077.1
Benguet		47.7	422.3	1,734.9	931.0	High	Lower Bound	-10.7	42.6	-14.2	362.5	-24.3	1,312.8	-16.1	782.0
-						Emission	Median	5.1	50.1	-5.5	399.2	-10.8	1,548.4	-5.2	882.9
						(RCP8.5)	Upper Bound	38.0	65.8	34.5	567.9	18.5	2,055.2	26.3	1,176.5
						Moderate	Lower Bound	4.8	107.6	-7.2	297.8	-35.8	687.1	~14.6	619.1
	ш					Emission	Median	7.4	110.2	-0.5	319.3	-16.4	895.7	-8.1	666.1
Ifugao	6	102.6	321.0	1,071.1	724.9	(RCP4.5)	Upper Bound	29.8	133.1	18.3	379.6	2.4	1,096.5	1.9	738.6
Ifu	2	102.0	521.0	1,071.1	724.5	High	Lower Bound	-5.2	97.3	-8.1	294.9	-27.8	773.0	-8.9	660.1
Ш	П					Emission	Median	7.9	110.7	1.3	325.0	-2,0	1,049.2	0.0	724.9
	_					(RCP8.5)	Upper Bound	33.5	137.0	30.5	419.1	13.6	1,217.2	11.6	809.3
						Moderate	Lower Bound	4.8	96.7	-3.7	219.6	-36.5	567.0	-13.8	596.7
	,					Emission (PCP4.5)	Median	7.3	99.0	3.3	235.6	-20.6	708.2	-5.0	657.3
Kalinga	2	92.3	228.0	892.3	691.9	(RCP4.5)	Upper Bound	27.0	117.2	17.8	268.6	3.9	927.5	6.8	738.8
Ka		1123		6000	7,500	High	Lower Bound	-4.1	88.5	-12.6	199.4	-27.7	645.4	-10.5	619.6
						(RCP8.5)	Median	9.7	101.3	0.2	228.4	-7.9	821.9	-0.4	688.9
	+					(40.0)	Upper Bound	22.6	113.2	24.0	282.7	9,9	981.1	11.6	772.1
Joe	3					Moderate	Lower Bound	3.9	77.7	-7.9	264.1	-21.0	885.3	-18.0	573.1
ivo						(RCP4.5)	Median	10.5	82.6	-2.7	278.9	-14.8	955.6	-8.9	636.8
n P		74.8	286.8	1,121.1	699.2	(1014.3)	Upper Bound	28.8	96.3	16.3	333.6	3.8	1,163.9	6.2	742.5
Mountain Province					1	High	Lower Bound	-5.8	70.5	-15.4	242.6	-20.0	896.9	-10.1	628.9
1 3	3					(RCP8.5)	Median Upper Bound	10.1 21.5	82.3 90.9	1.9	292.2 356.2	-3.0 16.9	1,087.9	0.5 13.8	702.7 795.7

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile



Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

I I		Observed (1971-2000)					Projected (2036-2065)								
-	90	200		ACC	SON	Scenario	Range*	DJF (Dec-Jan-Feb)		MAM (Mar-Apr-May)		JJA (Jun-Jul-Aug)		SON (Sep-Oct-Nov)	
Region	Province	DJF	MAM					Percent change	Projected value	Percent change	Projected value	Percent change	Projected value	Percent change	Projected value
П						Moderate	Lower Bound	-0.5	282.9	+8.2	190.7	-23.6	411.1	-6.5	778.4
۱	Cagayan	284.4			832.1	Emission	Median	7.4	305.4	2.3	212,5	-18.2	440.3	3.7	863.2
1			207.7	538.4		(RCP4.5)	Upper Bound	43.1	407.0	16,2	241.3	1.8	548.3	17.9	981.2
١		1000000				High	Lower Bound	-5.8	267.8	-15.2	176.1	-24.2	408.1	-5.8	783.7
ı						(RCP8.5)	Median	7.9	306.8 359.9	3.3 16.8	214.6 242.6	-2.2 17.4	526.7 632.2	1.3	842.9 980.1
ł						(1101010)	Upper Bound Lower Bound	3.5	426.4	1.1	328.7	-27.7	383.6	-4.4	829.0
1	Isabela	412.2	325.0	530.8	867.0	Moderate Emission (RCP4.5)	Median	11.5	459.7	10.3	358.5	-17.2	439.7	3.0	892.9
١							Upper Bound	48.4	611.7	23.5	401.3	0.3	532.4	11.9	970.1
1						High Emission (RCP8.5)	Lower Bound	-1.5	405.9	-6.1	305.3	-24.2	402.5	-2.3	847.1
,							Median	12.4	463.5	7.0	347.9	-2.6	516.8	11.0	962.7
5							Upper Bound	34.8	555.5	17.3	381.4	25.7	667.1	16.2	1,007.2
Kegion	Nueva Vizcaya	180.9	416.8	1,149.8	880.5	Moderate	Lower Bound	6.3	192.2	-0.1	416.2	-27.4	835.2	-14.9	749.5
٦						Emission (RCP4.5) High Emission (RCP8.5)	Median	9.4	197.9	5.6	440.1	-15.8	967.9	-5.5	832.2
							Upper Bound	40.2	253.6	29.1	538.2	-4.1	1,103.2	6.9	941.2
1							Lower Bound	-2.3	176.7	-6.5	389.5	-22.2	894.7	-8.6	804.5
ı							Median	8.6 39.9	196.5	30.9	422.6	-2.9 12.7	1,116.5	1.6	894.2
ł							Upper Bound Lower Bound	4.4	253.2 437.4	0.9	545.4 470.0	-27.3	1,295.9 564.8	-13.9	1,036.7 824.9
1	Quirino		465.9	776.4	957.9	Moderate Emission (RCP4.5)	Median	11.4	466.6	9.0	507.9	-16.9	645.1	-13.9	909.9
-		419.0					Upper Bound	57.3	659.0	31.7	613.5	-5.7	732.2	10.4	1,057.5
-							Lower Bound	-3.1	406.2	-8.5	426.1	-21.9	606.6	-10.6	856.4
-	0					High Emission (RCP8.5)	Median	9.4	458.3	1.5	473.0	-1.0	769.0	3.2	988.9
-							Upper Bound	52.1	637.4	26.5	589.6	17.7	913.9	17.5	1,125.3
7						Madarata	Lower Bound	1.5	624.7	1.6	555.0	-21.7	601.8	-9.7	1,039.6
-	Aurora	615.7	546.4	768.7	1,151.1	Moderate Emission	Median	8.5	667.9	6.8	583.7	-8.6	702.7	-2.4	1,123.8
-						(RCP4.5)	Upper Bound	52.0	936.0	22.5	669.2	-6.8	716.2	5.3	1,211.6
-						High Emission (RCP8.5)	Lower Bound	-6.1	578.2	-10.2	490.8	-16.6	641.1	-5.0	1,093.9
ı							Median	3.0	633.9	4.6	571.6	-3.4	742.6	2.8	1,183.6
1							Upper Bound	31.8	811.7	19.0	650.0	10.2	847.1	11.1	1,278.6
	Bataan	71.7	368.7	1,326.2	872.6	Moderate Emission (RCP4.5)	Lower Bound	1.0	72.4	-3.3	356.7	-25.5	988.1	-10.0	785.0
П							Median	12.7	80.8	12.2	413.7	-17.7	1,091.1	-1.9	856.0
1						High Emission (RCP8.5)	Upper Bound	59.2	114.2	18.7	437.8	-1.0	1,313.3	9.3	953.6
1							Lower Bound	-2.1	70.2	-9.7	333.0	-19.0	1,074.7	-6.2	818.6
							Median	20.2	86.2	4.6	385.6	-1.9	1,301.3	3.1	899.3
H						(((0.5)	Upper Bound Lower Bound	38.2	99.1	25.3	461.9 289.0	-23.2	1,490.1 799.4	18.3 -8.6	769.6
П	Bulacan	212.4	288.9	1,041.4	842.1	Moderate Emission (RCP4.5) High Emission (RCP8.5)	Median	18.0	250.6	12.5	325.1	-10.7	929.7	-5.8	793.5
1							Upper Bound	61.5	343.1	23.8	357.8	1.4	1,056.4	11.6	939.5
-							Lower Bound	-2.0	208.2	-6.8	269.1	-18.3	850.7	-9.2	764.5
							Median	15.4	245.1	3.3	298.4	-5.0	989.2	1,2	852.4
							Upper Bound	48.1	314,5	22.9	355.1	9.6	1,141.7	18.3	995.9
						Moderate	Lower Bound	4.3	161.9	0.6	318.3	-20.0	795.5	-9.5	674.2
m	Nueva Ecija			995.0	745.0	Emission	Median	14.6	177.9	10.9	351.0	-16.4	832.2	-5.8	701.5
Region		155.2	316.5			(RCP4.5)	Upper Bound	55.0	240.6	24.1	392.7	0.9	1,004.0	16.6	868.5
Reg			-	2000		High Emission (RCP8.S)	Lower Bound	-6.5	145.1	-6.9	294.7	-18.6	809.9	-10.8	664.5
							Median	10.7	171.8	3,8	328.6	-7.9	916.5	2.1	760.8
1		-					Upper Bound	45.3	225.5	20,4	381.0	10.7	1,101.7	16.6	868.4
	Pampanga			1,030.4	785.2	Moderate Emission (RCP4.5)	Lower Bound	4.9	126.8 143.3	9.0	333.5	-22.8 -12.3	795.0	-8.0 -1.9	722.5 769.9
		200					Median Upper Bound	18.6 75.8	212.4	19.7	349.6 383.9	4.8	903.3	7.2	841.9
		120.8	320.6				Lower Bound	-4.4	115.5	-9.1	291.4	-12.0	907.0	-5.2	744.7
						High Emission (RCP8.5)	Median	18.5	143.1	6.5	341.4	-0.8	1,022.4	-1.2	776.0
							Upper Bound	80.7	218.3	18.5	380.1	10.0	1,133.6	23.8	971.9
1	Tarlac	43.4	265.4	1,193.5	644.3	Moderate Emission	Lower Bound	3.3	44.8	3.2	273.8	-18.3	975.4	-8.3	591.0
							Median	7.9	46.8	8.1	287.0	-8.3	1,094.7	-4.7	613.9
						(RCP4.5) High Emission (RCP8.5)	Upper Bound	49.3	64.8	21.2	321.8	0.8	1,203.5	18.1	761.0
							Lower Bound	-5.7	40.9	-11.7	234.4	-15.9	1,004.1	-3.3	623.2
							Median	9.9	47.7	5.0	278.6	-5.1	1,133.0	1.8	655.8
							Upper Bound	33.4	57.9	24.8	331.2	8.4	1,293.2	17.6	757.7
Í		11 71				Moderate	Lower Bound	-1.5	40.3	-0.4	366.5	-22.6	1,389.0	-9.7	787.4
	Zambales	40.9	368.0	1,793.9	872.0	Emission	Median	11.3	45.5	7.4	395.3	-14.4	1,536.1	-2.5	850.3
						(RCP4.5)	Upper Bound	51.9	62.1	18.5	436.2	0.9	1,809.6	14.0	994.1
						High Emission (RCP8.5)	Lower Bound	-5.9	38.5	-16.4	307.8	-16.5	1,497.3	-3.9	838.0
- 1							Median Upper Bound	9.2	44.7 54.2	1.6	373.9 439.5	-4.7 12.7	1,710.1 2,021.2	4.6 14.9	911.9

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

		Observed (1971-2000)							Projected (2036-2065)						
=	age					Scenario	Range*	DJF (Dec-Jan-Feb)		MAM (Mar-Apr-May)		JJA (Jun-Jul-Aug)		SON (Sep-Oct-Nov)	
Region	Province	DJF	MAM	JJA	SON			Percent change	Projected value	Percent change	Projected value	Percent change	Projected value	Percent change	Projected value
						Moderate	Lower Bound	7.5	248.2	-0,1	280.1	-27.7	619.4	-11.4	661.6
П	Se				746.4	Emission (RCP4.5)	Median	14.9	265.5	1.7	285,1	-15.2	726.1	-3.6	719.7
1	Batangas	231.0	280.4	856.5		(RCP4.5)	Upper Bound	56.1	360.7	15.8	324.8	1.0	865.1	8.2	807.2
	Bat					High Emission (RCP8.5)	Lower Bound	1.4	234.2	-11.4	248.3	-23.5	655.6	-9.7	673.7
							Median Upper Bound	8.1 37.7	249.8 318.2	0.5 25.1	281.8 350.7	-8.9 10.2	779.9 944.0	10.0	749.5 821.2
1							Lower Bound	8.6	135.6	0.6	244.3	-26.7	722.4	-7.7	534.3
	Cavite	124.9	242.8	985.7	579.0	Moderate Emission (RCP4.5)	Median	12.5	140.5	6.9	259.6	-18.0	808.1	-4.1	555.4
							Upper Bound	55.7	194.5	17.9	286.2	9.4	1,078.6	6.7	618.0
١						High Emission (RCP8.5)	Lower Bound	7.8	134.7	-11.0	216.1	-22.8	760.8	-4.3	553.9
١							Median	12.8	140.9	3.7	251.8	-10.4	883.2	0.9	584.2
							Upper Bound	35.9	169.7	33.1	323.1	11.7	1,100.6	10.3	638.9
		629.2			1	Moderate	Lower Bound	4.2	655.9	-2.1	378.5	-22.7	653.4	-9.0	970.3
4-A	Laguna				1,066.5	Emission (PCD4 E)	Median	10.2	693.2	12.6	435.4	-14.3	724.0	-5.8	1,004.7
6			386.8	845.0		(RCP4.5) High Emission (RCP8.5)	Upper Bound	43.9	905.2	24.8	482.6	-2.1	827.3	5.7	1,127.0
Region							Lower Bound	2.3	643.7	-14.4	331.0	-20.9	668.8	-10.8	951.0
٦							Median	13.8	716.3 836.4	-0.1 28.3	386.3 496.2	-11.0 7.5	752.0	1.4	1,081.1
1							Upper Bound Lower Bound	32.9 4.7	866.2	6.2	406.4	-22.0	908.8 522.4	-9.3	1,178.9
	Quezon				1,229.3	Moderate Emission (RCP4.5)	Median	8.4	897.2	9.3	418.4	-17.2	554.9	1.6	1,249.0
		827.7		670.0			Upper Bound	31.6	1,089.2	18.9	455.1	5.3	705.8	7.6	1,322.5
			382.7			OU-L	Lower Bound	-0.8	820.8	-4.5	365.6	-16.9	557.0	-7.0	1,143.2
						High Emission (RCP8.5)	Median	13.9	942.6	7.3	410.6	-6.3	627.8	2.5	1,260.3
							Upper Bound	32.4	1,095.7	19.6	457.9	5.6	707.8	14.0	1,400.9
1						Moderate	Lower Bound	0.0	262.4	-2.8	234.8	-27.3	728.0	-14.9	699.5
		262,4				Emission	Median	7.8	282.8	12.9	272.7	-20.0	800.7	-9.5	743.7
	Rizal		241.5	1,001.3	821.8	(RCP4.5)	Upper Bound	51.5	397.4	25.6	303.4	-1.7	983.9	12.7	926.5
						High Emission (RCP8.5)	Lower Bound	3.6	271.8	-14.2	207.2	-25.4	747.4	-13.2	713.6
4							Median	15.0	301.7	-1.1	238.8	-11.6	885.5	0.9	829.6
4						(ACFO.3)	Upper Bound	52.2	399.4	17.1	282.7	11.7	1,118.9	19.6	983.1
	Occidental Mindoro	159.5	265.9	1,091.2	762.6	Moderate	Lower Bound	-1.7	156.8	-5.0	252.6	-25.1	817.5	-19.5	613.8
						(RCP4.5)	Median Upper Bound	13.2 52.3	180.6 242.9	2.6 12.7	272.8 299.6	-20.6 3.6	866.8 1,130.2	-3.0 4.2	740.0 794.3
						High Emission (RCP8.5)	Lower Bound	-1.4	157.3	-18.3	217.2	-28.1	784.2	-18.1	624.9
							Median	9.5	174.6	-3.3	257.2	-13.0	949.8	-3.6	735.1
	000						Upper Bound	25.9	200.7	23.3	327.9	11.6	1,217.6	7.9	822.5
1		260.3	269.3	894.3	791.2	Moderate Emission (RCP4.5) High Emission (RCP8.5)	Lower Bound	0.0	260.2	3.6	279.1	-24.9	671.2	-17.3	654.6
П	Oriental Mindoro						Median	10.8	288.5	5.6	284.3	-18.8	726.2	-7.7	729.9
П							Upper Bound	31.9	343.3	12.3	302.5	-0.4	890.8	5.2	832,3
	nta						Lower Bound	-2.7	253,3	-9.5	243.7	-27.2	650.7	-13.9	681.2
4-B	ore						Median	13.9	296.6	5.7	284.6	-12.9	778.9	-6.8	737.5
5	-						Upper Bound	27.9	332.8	9.2	294.1	9.5	979.6	4.9	829.8
Region	an				640.6	Moderate Emission (RCP4.5) High Emission (RCP8.5)	Lower Bound	-9.9	91.8	-10.6	169.2	-25.7	581.1	-14.3	548.7
-							Median	0.0	101.8	-7.4	175.3	-12.0	688,2	-8.1	588.9
	Palawan	101.8	189.3	781.7			Upper Bound Lower Bound	26.8 -17.8	129.1 83.6	10.3 -16.2	208.9 158.6	-25.0	789.3 586.2	10.2 -22.5	705.7 496.5
	P						Median	4.3	106.2	-3.6	182.4	-5.6	737.8	-6.2	601.0
							Upper Bound	14.2	116.3	9.2	206.7	12.9	882.5	11.1	711.5
						Moderate	Lower Bound	1.7	363.2	-5.2	212.3	-26.0	483.2	-13.3	674.8
	Rombion	357.0	224.0	653.0	778.0	Emission	Median	10.6	395.0	5.0	235.2	-21.0	515.8	1.5	789.4
						(RCP4.5)	Upper Bound	37.1	489.6	31.7	295.0	4.3	681.2	13.6	883.6
	Rom					High Emission	Lower Bound	-3.6	344.0	-15.8	188.5	-35.3	422.3	-25.6	578.5
							Median	10.8	395.5	6.4	238.4	-6.6	609.6	-4.6	742.3
	214					(RCP8.5)	Upper Bound	26.0	449.7	23.7	277.1	15.1	751.5	20.5	937.4
	Metro Manila	107.5		1,170.2	758.7	Moderate Emission (RCP4.5)	Lower Bound	-0.1	107.3	0.7	199.8	-21.3	920.8	-10.8	676.4
2							Median	17.7	126.5	6.9	212.2	-10.1	1,051.6	-6.0	713.5
NCR			198.5			High Emission (RCP8.5)	Upper Bound	55.5	167.1	25.7	246.9	-0.4	1,165.2	7.7	817.2
	letr						Lower Bound	2.7	110.4	-7.2 4.8	184.2	-17.0 -6.1	970.9	-8.0 3.9	698.1 788.3
	2						Median Upper Bound	53.4	164.9	19.8	208.1	7.7	1,098.5	19.9	909.4
-							Lower Bound	-3.5	713.9	-2.4	377.6	-25.5	525.6	-12.1	827.2
	Albay	739.8	386.9	705.8	941,3	Moderate Emission (RCP4.5)	Median	13.5	839.4	2.8	397.9	-12.8	615.7	-2.6	916.6
S IIC							Upper Bound	38.7	1,026.3	20.1	464.6	-2.5	688.1	8.7	1,023.3
Region 5							Lower Bound	-4.8	704.6	-0.2	386.1	-25.2	527.9	-11.0	837.9
~						Emission	Median	14.2	844.5	5.5	408.3	-4.7	672.8	1.4	954.9
						(RCP8.5)	Upper Bound	40.3	1,037.7	18.1	456.8	11.9	789.8	8.1	1,017.4

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

		Obs	served (1971-20	00)				Pro	jected (20	036-2065)				
	Joe	The same of				1		DJF (Dec	-Jan-Feb)	MAM (Mar	r-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	MAM	JJA	SON	Scenario	Range*	Percent change	Projected value	Percent change	Projected value	Percent change	Projected value	Percent change	Projected value
ï	te					Moderate	Lower Bound	2.2	1,051.9	6.8	425.4	-21.6	443,5	-17.1	1,066.2
	Camarines Norte		100			Emission (RCP4.5)	Median	7.9	1,111.0	11.7	445.0	-18.1	463.5	-1.3	1,269.3
	nes	1,029.6	398.5	565.6	1,285.7	(NCF4.3)	Upper Bound	35.4	1,394.0	14.3	455.4	3.7	586.4	7.5	1,381.5
	mar	1	100		7	High Emission	Lower Bound Median	-1.9 19.9	1,009.9	-6.4 7.3	373.1 427.7	-21.6 -7.5	443.3 523.4	-8.4	1,177.3
	S		-			(RCP8.5)	Upper Bound	31.0	1,348.5	18.9	473.7	8.6	614.1	14.9	1,477.7
P	-						Lower Bound	-0.7	662.3	2.1	354.6	-25.5	476.6	-14.3	882.7
	Sur					Moderate Emission	Median	13.3	755.3	5.5	366.5	-17.6	527.1	-2.5	1,003.8
	nes	666.8	347.4	639.6	1,029.4	(RCP4.5)	Upper Bound	37.7	918.4	15.6	401.6	1.0	645.8	2.9	1,059.1
Н	Camarines	000.8	347.4	039.0	1,029.4	High	Lower Bound	-3.0	647.0	-3.1	336.5	-25.1	479.3	-8.3	944.3
	S					Emission (RCP8.5)	Median	17.1	780.9	6.8	371.1	-8.5	585.1	-0.2	1,027.3
ļ,	111					(NCF0.5)	Upper Bound	27.8	852.0	15.8	402.5	11.3	712.2	6.2	1,092.9
J	- CO					Moderate	Lower Bound Median	-2.1 13.6	1,052.5	-0.9 9.7	507.9 562.4	-27.9 -17.6	465.9 532.8	-9.8 -6.6	1,081.4
n 5	Catanduanes		1,300			(RCP4.5)	Upper Bound	33.5	1,435.1	27.7	654.6	3.9	671.4	0.4	1,204.1
Region	npu	1,075.4	512.7	646.3	1,199.5		Lower Bound	0.5	1,080.8	-1.7	504.2	-30.3	450.3	-17.6	988.9
ă	Cata					High Emission	Median	15.0	1,237.1	4.5	535.5	-10.2	580.2	0.6	1,206.3
						(RCP8.5)	Upper Bound	28.6	1,383.1	19.9	615.0	7.9	697.2	9.7	1,315.7
						Moderate	Lower Bound	3.6	528.6	0.9	253.0	-18.5	463.8	-11.0	658.1
	-61					Emission	Median	10.5	563.5	10.6	277.3	-12.2	500.1	-0.4	736.1
	Masbate	510.2	250.7	569.4	739.3	(RCP4.5)	Upper Bound	40.7	718.1	18.7	297.5	2.9	586.0	8.1	798.9
	Mas	310.2	230.7	303.1	7.55.5	High	Lower Bound	-6.1	479.3	-4.3	240.0	-21.8	445.1	-16.9	614.0
						(RCP8.5)	Median	11.1	566.8	7.1	268.4	-7.3	527.7	-3.8	711.0
b	_				_	(1101013)	Upper Bound	28.8	656.9 970.8	17.1 -0.2	293.7 427.2	-20.7	608.6	16.3 -5.1	859.5 923.6
						Moderate Emission	Lower Bound Median	10.2	1,055.6	6.9	457.6	-7.9	524.0 608.4	-1.8	956.6
	gon	77.0	F. (2. 7.7.7			(RCP4.5)	Upper Bound	31.6	1,260.9	18.4	506.8	2.1	674.5	1.1	983.9
	Sorsogon	958.1	427.9	660.4	973.6	1001	Lower Bound	-2.2	936.6	4.5	447.1	-23.6	504.7	-14.4	833.0
	Š				- 1	High Emission	Median	13.1	1,083.4	7.3	459.0	-4.0	633.9	-1.2	962.2
	11					(RCP8.5)	Upper Bound	36.0	1,303.0	18.3	506.3	5.8	699.0	9.4	1,065.4
						Moderate	Lower Bound	-0.3	429.9	-6.8	300.8	-22.0	672.5	-14.0	759.6
						Emission	Median	4.6	450.9	0.2	323.5	-10.5	771.9	-0.8	876.9
	Aklan	431.2	322.7	862.5	883.7	(RCP4.5)	Upper Bound	29.7	559.4	7.7	347.7	0.5	866.7	6.3	939.2
	A		20.00		100	High	Lower Bound	-6.2	404.5	-9.7	291.4	-20.3	687.0	-21.5	693.5
						(RCP8.5)	Median	11.0	478.7	-3.4	311.6	-9.4	781.2	-4.8	841.2
r							Upper Bound Lower Bound	-11.9	527.6 262.5	10.7 -21.6	357.3 225.9	-32.2	903.6 675.2	5.5 -27.4	932.4 610.7
						Moderate Emission	Median	-2.1	291.6	-14.6	245.9	-10.5	890.8	-14.8	717.3
	ane	2022		1000		(RCP4.5)	Upper Bound	34.2	399.8	7.5	309.6	1.9	1,014.3	-0.9	833.5
	Antique	297.9	288.0	995.3	841.4	High	Lower Bound	-20.3	237.3	-31.8	196.3	-38.0	617.3	-30.6	584.1
						Emission	Median	6.9	318.5	-13.0	250.7	-8.8	907.9	-8.6	768.9
						(RCP8.5)	Upper Bound	20.3	358,3	4,3	300.4	9.0	1,085.1	2,2	859.6
						Moderate	Lower Bound	3.0	483.7	-4.0	328.2	-17.8	669.4	-15.5	751.7
						(RCP4.5)	Median	6.9	502.1	2,0	348.7	-6.9	757.7	-5.3	842,4
	Capiz	469.7	342.0	814.2	889.1		Upper Bound	33.9	628.8 457.0	13.5	388.1	8.3 -18.2	881.8	-21.5	928.4 697.5
1	U					High Emission	Lower Bound Median	11.8	525.3	-6.4 0.6	320.1 344.1	-18.2	665.9 735.8	-6.8	828.5
on 6						(RCP8.5)	Upper Bound	25.7	590.4	12.9	386.2	8.0	879.4	5.4	937.4
Region						Moderate	Lower Bound	4.7	237.8	-1.0	231.2	-14.6	720.7	-15.1	594.2
ď			144			Moderate Emission	Median	13.6	257.9	3.4	241.4	-6.6	788.2	-13.0	608.5
	Guimaras	227.0	233.0	843.0	699.0	(RCP4.5)	Upper Bound	24.6	282.8	12.9	263.7	11.4	939.6	12.2	785.2
	Suin	227.0	233.0	043.0	099.0	High	Lower Bound	-4.2	217.5	-10,8	208.4	-18.7	686.0	-17.9	574.7
	9					Emission	Median	7.7	244.5	3.2	241.0	-3.9	810.8	-10.1	629.0
į,	-	-				(RCP8.5)	Upper Bound	27.9	290.5	18.0	275.5	14.0	961.5	9.3	764.5
						Moderate	Lower Bound	-4.4	310.4	-9.3	263.4	-22.1	726.9	-20.0	662.6
		Jan Hill				(RCP4.5)	Median	6.5	346.0	4.0	302.1	-5.8	878.4	-11.2	735.9
	Ilollo	324.8	290.6	932.8	828.3		Upper Bound Lower Bound	26.9 -9.8	412.1 293.0	-15.4	324.5 246.0	7.1	998.6 667.9	7.9	894.1 639.9
			-			High Emission	Median	8.7	353.2	-2.1	284.6	-4.0	895.0	-11.1	736.4
						(RCP8.5)	Upper Bound	28.3	416.8	9.5	318.2	10.3	1,028.6	3.4	856.2
ľ	7	-				Moderate	Lower Bound	-12.0	206.7	-12.7	247.1	-20.6	714.0	-21.5	615.3
	ents		10.1			Moderate Emission	Median	1.8	239.1	0.1	283.2	-6.5	841.4	-14.3	671.7
	Negros Occidental	234.9	283.0	899.6	784.0	(RCP4.5)	Upper Bound	25.2	294.1	11.0	314.0	3.1	927.4	1.5	795.5
Ш	0 50	234.9	203.0	0,666	704.0	High	Lower Bound	-17.3	194.3	-18.6	230.4	-21.2	708.9	-27,9	565.4
	legn					Emission	Median	6.6	250.3	-7.0	263.3	-5.2	853.1	-13.7	676.6
	2					(RCP8.5)	Upper Bound	30.0	305.3	2.8	290.9	5.9	952.3	1.2	793.4

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

		Obs	served (1971-20	00)				Pro	jected (20	36-2065)	(
-	Joe	1				1	4-3-7	DJF (Dec	-Jan-Feb)	MAM (Mar	-Apr-May)	JJA (Jun-	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	MAM	JJA	SON	Scenario	Range*	Percent change	Projected value						
						Moderate	Lower Bound	-13.5	325.3	-2.4	204.5	-16.8	343.5	-21.2	405.7
						Emission	Median	3.0	387.5	3.9	217.7	-6.5	386.1	-12.2	451.9
	Bohol	376.1	209.6	412.9	514.5	(RCP4.5)	Upper Bound	29.7	487.8		234.5	13.8	470.1	4.8	539.1
	Bo	370(1	203.0	144.5	54115	High	Lower Bound	-17.1	311.9		195.5	-22.1	321.7	-27.9	370.8
						Emission (RCP8.5)	Median	1.5	381.8	1.8	213.3	-2.3	403.3	-12.3	451.4
,						(KCPO.5)	Upper Bound	21.5	457.1	8.5	227.5	13.4	468.2	6.3	546.8
П						Moderate	Lower Bound	-7.1	301.1	-3.7	219.8	-12.4	521.3	-17.7	499.9
17	- 4		1623			(RCP4.5)	Median	3.9	336.6		243.4	-4.9	565.7	-6.0	570.7
Region	Cebu	324.0	228.3	595.1	607.4	1000.009	Upper Bound	31.7 -15.4	426.9 274.1	9.2	249.4 221.1	1.0 -15.8	601.2 501.1	1.5 -21.3	616.4 478.0
Re	ŭ					High Emission	Lower Bound Median	6.2	344.1	2.1	233.1	-5.4	562.8	-11.5	537.5
П						(RCP8.5)	Upper Bound	22.4	396.7	10.2	251.5	3.6	616.6	2.7	623.5
P							Lower Bound	-12.6	197.4	-8.5	206.8	-20.4	509.3	-18.2	520.8
П	ital					Moderate Emission	Median	2.3	231.0	-4.1	216.8	-7.8	589.5	-14.1	547.3
	Negros Orienta	1000	2000	week.	a Destruction	(RCP4.5)	Upper Bound	23.5	278.9	8.4	245.1	8.6	694.2	-1.6	626.8
	0 50	225.8	226.0	639.5	636.9	CUSA	Lower Bound	-14.9	192.2	-22.7	174.6	-20.8	506.4	-26.4	468.8
	egr					High Emission	Median	4.1	235.1	-7.0	210.2	-1.2	632.0	-14.5	544.7
	Z					(RCP8.5)	Upper Bound	20.2	271.4	7.3	242.5	7.6	688.1	3.8	661.2
						Moderate	Lower Bound	-4.2	945.9	-2.6	452.0	-6.7	522.3	-20.1	696.5
Ш	Samar					Emission	Median	1.3	999.5	1.8	472.3	-1.1	553.8	-16.6	726.7
Ш	Sa	987.0	464.1	559.8	871.4	(RCP4,5)	Upper Bound	45.0	1,431.4	17.4	544.9	8.3	606.2	-3.8	838.0
Ш	Eastern	907.0	404.1	339.0	0/1.4	High	Lower Bound	-7.9	909.2	5.8	490.9	-14.0	481.5	-22.0	680.1
Ш	Eas					Emission	Median	13.7	1,122.2	1.1	469.4	-0.6	556.6	-9.3	790.1
Н						(RCP8,5)	Upper Bound	43.6	1,417.8		524.4	9.7	614.4	4.7	912.1
П						Moderate	Lower Bound	-7.5	637.6		323.4	-18.5	463.6	-24.3	549.3
Ш						(RCP4.5)	Median	0.4	692.5		377.3	-1.1	562.3	-15.5	612.9
Ш	Leyte	689.5	342.0	568.7	725.5	(0014:5)	Upper Bound	35.2	931.9		382.9	8.1	614.7	1.0	733.0
Ш	2	777	1		1 200	High	Lower Bound	-15.2	584.9		318.0	-19.8	456.0	-26.4	534.0
П						(RCP8.5)	Median	9.2	752.9	-1.2	338.0	-3.5	548.7	-9.8	654.7
					_	(merolo)	Upper Bound	31.9	909.8		357.1	6.6	606.2	6.5	772.6
J	Jar.					Moderate	Lower Bound	7.2	1,154.1	-0.7 2.2	458.8	-14.9 -7.3	482.4	-14.4 -8.4	839.7
n 8	Samar	Um)	100	40.0		(RCP4.5)	Median Upper Bound	45.5	1,210.4		472.5 546.7	3.4	525.6 586.1	3.7	898.8 1,017.5
Region		1,128.9	462.2	566.8	981.4		Lower Bound	-4.8	1,074.2	0.0	462.1	-19.5	456.5	-21.1	774.8
æ	Northern		12 11			High Emission	Median	14.8	1,295.8	-	478.5	-4.0	544.3	-3.2	949.7
П	ž					(RCP8.5)	Upper Bound	43.0	1,614.8		524.0	6.3	602.7	7.3	1,052.7
	1					A Paral Control	Lower Bound	-4.8	846.4	-4.7	416.7	-18.9	486.4	-18.8	713.7
Ш						Moderate Emission	Median	4.1	926.4	5.5	461.0	-8.1	551.0	-10.9	783.4
Ш	nar	000 5	427.0	500.0	070 4	(RCP4.5)	Upper Bound	45.8	1,297.0	20.6	526.9	7.5	644.5	1.9	896.2
Ш	Samar	889.5	437.0	599.8	879.4	High	Lower Bound	-10.5	796.3	-3.0	423.9	-23.8	457.3	-21.9	687.2
Ш						Emission	Median	8.3	963.2	-0.4	435.2	-3.2	580.3	-5.5	831.0
Ш						(RCP8.5)	Upper Bound	34.4	1,195.6	10,2	481.7	6.6	639.5	2.2	898.9
П	- CU					Moderate	Lower Bound	-10.2	735.4	-3.1	350.9	-16.9	424.1	-20.2	555,1
Ш	eyt					Emission	Median	-2.7	796.5	4.5	378.6	-3,4	493.5	-10.6	622,1
	uthern Leyte	818.6	362.2	510.6	695.6	(RCP4.5)	Upper Bound	27.2	1,041.4	10.4	399.9	7.8	550.2	3,3	718.8
Ш	ithe	BILLION,	7,5	20000		High	Lower Bound	-12.9	712.8		348.7	-10,4	457.3	-25.6	517.4
	Sol					(RCP8.5)	Median	13.3	927.4		372.8	-3.4	493.4	-11.0	619.4
Н			-			(10.0.0)	Upper Bound	35.8			413.7	12,2	573.1	9.5	761.9
П	del					Moderate	Lower Bound	-14.9	276.1	-19.2	226.0	-10.3	537.3	-21.1	566.3
Ш	ga d					(RCP4.5)	Median	-3.7	312.5		255.9	-2.0	587.2	-4.4	686.6
Ш	lort	324.5	279.7	599.1	718.1	(ittel 115)	Upper Bound	25.4	407.0		297.0	14.9	688.1	6.6	765.3
Ш	Zamboanga		5000			High	Lower Bound	-12.4	284.2		234.8	-10.4	537.0	-18.7	583.9
	Za		-			(RCP8.5)	Median Upper Bound	1.9	330.6 369.3		272.8 302.5	4.9 12.2	628.6 672.0	-0.5 10.8	714.2 795.4
							Lower Bound	-8.2	270.4		238.5	-11.9	523.0	-23.6	506.6
	del					Moderate Emission	Median	-4.1	282.4	-7.6		-2.5	579.2	-6.0	623.4
6 u	- Jaa			3.00		(RCP4.5)	Upper Bound	23.1	362.6		319.0	9.4	649.5	8.8	721.7
Region	Sul	294.5	298.7	593.8	663.2	inc.	Lower Bound	-18.1	241.3		274.2	-11.8	523.9	-13.3	574.7
R	Zamboanga del Sur	7 7	4		5.71	High Emission	Median	4.4	307.6		288.9	2.0	605.5	-0.6	659.0
	7	11-11				(RCP8.5)	Upper Bound	13.8	335.2	16.3	347.4	16.7	693.0	6.2	704.2
1						Market 1	Lower Bound	-14.4	243.1	-9.8	262.0	-12.0	525.4	-21.5	528.9
	ag		100			Moderate Emission	Median	-6.6	265.4		283.1	-3.9	573.7	-5.4	637.9
П	Zamboanga Sibugay	2011		2000	200	(RCP4.5)	Upper Bound	12.3	319.1	7.0	310.9	8.3	646.9	3.3	696.1
	mpc	284.1	290.5	597.2	674.1	Mich	Lower Bound	-11.6	251.2		236.9	-15.4	505.3	-14.8	574.0
	Za		10.11		1	High Emission	Median	-0.2	283.6		293.5	2.3	610.8	-6.0	633.6
						(RCP8.5)	Upper Bound	8.4	308.1	7.5	312.2	13.4	677.1	7.0	721.5

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

		Obs	served (1971-20	00)	21	,		Pro	jected (20	36-2065)				
	ace	405				7		DJF (Dec-	Jan-Feb)	MAM (Mar	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	МАМ	JJA	SON	Scenario	Range*	Percent change	Projected value						
ľ						Moderate	Lower Bound	-18.9	267.4	-21.0	265.0	-27.3	475.4	-41.1	329.8
	uc	1	44.0			Emission (RCP4.5)	Median	-3.2	319.2	-9.7	303.1	-16.3	547.4	-26.5	411.2
	Bukidnon	329.7	335.6	653.8	559.5	(1001-110)	Upper Bound	17.3	386.8	12.8	378.5	12.6	736.5	-7.7	516.2
	Bul		100.0	4 1		High Emission	Lower Bound Median	-18.4 -7.4	269.1 305.3	-14.5 -6.7	287.0 313.1	-19.6 -4.4	525.9 625.1	-22.4 -10.9	434.1 498.6
						(RCP8.5)	Upper Bound	25.7	414.4	5.4	353.6	17.0	765.0	10.1	616.0
-	7					GARDINE.	Lower Bound	-9.5	305.4	-22.9	270.1	-8.5	606.4	-30.5	431.9
	Norte					Moderate Emission	Median	-2.4	329.3	-7.5	324.2	-1.8	650.5	-8.4	569.0
	del N	337.5	350.3	662.5	621.1	(RCP4.5)	Upper Bound	21.4	409.6	87.3	656.2	69.4	1,122.0	61.4	1,002.2
	go d	237.5	550.5	002.5	021.1	High	Lower Bound	-10.0	303.6	-16.0	294.3	-10.0	596.5	-15.4	525.4
10	Lanao					(RCP8.5)	Median	1.3	342.0	1.2	354.4	1.9	675.0	-2.3	606.9
Region						(Ner ols)	Upper Bound Lower Bound	13.2 -5.5	382.0 370.3	27.6 -11.0	447.1 288.0	19.7 -11.8	793.0 558.1	9.8	682.2 583.8
Rec	Misamis Occidental					Moderate Emission	Median	1.1	396.2	-5.6	305.2	3.5	655.2	-19.6	718.2
	cide	5000	12.00	0.000	1700	(RCP4.5)	Upper Bound	42.3	557.8	10.1	356.1	16.8	739.6	14.7	835.6
	s Oc	392.1	323.4	633.1	728.3	High	Lower Bound	-6.6	366.2	-7.2	300.1	-8.1	581.5	-6.4	681.4
	sam					Emission	Median	8.2	424.2	3.6	335.0	3.2	653.1	-1.0	720.9
	Σ					(RCP8.5)	Upper Bound	27.9	501.4	14.9	371.6	21.1	766.7	9.0	794.0
	- a					Moderate	Lower Bound	-21.4	347.6	-19.1	239.5	-22.2	478.8	-37.3	364.6
	ient					(RCP4,5)	Median	-6.8	412.3	-6.4	277.1	-14.1	529.0	-21.2	457.8
	Misamis Oriental	442.5	296.0	615.7	581.1	(NCF4,5)	Upper Bound	37.0	606.4	7.1	317.0	8.5	668.0	-0.7	576.8
	ami		-	-		High Emission	Lower Bound Median	-27.2	322.0 439.3	-20.0 -5.6	236.7 279.4	-15.9 -2.8	518.0 598.2	-33.2 -10.9	387.9 517.6
	N N					(RCP8,5)	Upper Bound	28.1	567.0	2.8	304.3	14.6	705.9	16.7	678.4
	>					District.	Lower Bound	-21.2	589.4	-17.8	459.8	-32.6	368.7	-37.5	366.6
	Compostella Valley					Moderate Emission	Median	-12.2	657.0	-8.6	511.0	-12.8	476.7	-22.7	453.3
	lla /	748.1	559.0	546.7	586.6	(RCP4.5)	Upper Bound	25.0	935.2	16.4	650.9	6.3	581.1	-4.7	558.8
	oste	740.1	559.0	340.7	300.0	High	Lower Bound	-23.8	570.1	-13.5	483.4	-22.6	423.1	-33.9	388.0
	ф					Emission	Median	-8.5	684.2	-5.1	530.3	-3.8	525.7	-4.2	561.9
þ	ŭ					(RCP8.5)	Upper Bound	15.3	862.9	8.4	606.1	16.4	636.1	7.6	631.3
	te					Moderate	Lower Bound	-14.4	545.1	-12.6	434.1	-22.1	417.5	-29.6	391.7
	Davao del Norte		1000			(RCP4.5)	Median Upper Bound	-3.3 20.3	616.1 766.6	-4.6 8.9	473,4 540.8	-9.4 6.2	485.1 568.6	-20.4 -7.2	442.8 516.0
	del	637.0	496.5	535.6	556.2		Lower Bound	-11.7	562.3	-10.6	443.9	-12.6	467.9	-22.3	432.4
	avac					High Emission	Median	-2.1	623.6	-2.4	484.5	-2.8	520.4	-11.0	495.2
	۵					(RCP8.5)	Upper Bound	24.2	791.3	11.6	553.9	13.0	605.5	10.5	614.5
ľ	77					Moderate	Lower Bound	-14.2	247.3	-8.8	316.7	-14.0	424.8	-24.1	335.9
11	Sur					Emission	Median	-3.6	277.9	-4.1	332.9	-1.8	485.3	-14.5	378.1
Ion	del	288.1	347.1	494.1	442.3	(RCP4.5)	Upper Bound	16.0	334.3	15.4	400.5	8.4	535.4	-3.7	425.8
Region	Davao del	State Street	200	100		High	Lower Bound	-8,4	263.9	-6.4 2.1	325.0	-7.1	459.2	-14.4	378.5
Ŧ	ā					(RCP8.5)	Median Upper Bound	-0.9 16.0	285.6 334.2	15.4	354.2 400.4	18.3	515.6 584.4	-10.1 8.2	397.8 478.4
	De 1					10.707.710	Lower Bound	-14.1	710.3	-3.1	593.1	-24.8	406.5	-20.5	476.4
	ental					Moderate Emission	Median	5.3	870.8	4.9	641.6	-1.8	530.4	-7.5	554.5
	Occidenta	827.3	C11.0	540.4	599.2	(RCP4.5)	Upper Bound	14.7	949.3	8.7	665.1	35.0	729.6	12.2	672.6
	0 0	027.3	611.8	540.4	599.2	High	Lower Bound	-6.2	776.1	-7.6	565.4	-19,3	436.3	-21.2	472.3
	Davao					Emission	Median	3,6	857.2	2.0	623.8	3,1	557.4	-2.5	584.5
	7					(RCP8.5)	Upper Bound	17.2	969.5	10.5	676.0	22.7	663.1	9.3	655.1
	70					Moderate	Lower Bound	-13.4	716.5	-8.3	561.2	-21.7	423.0	-27.8	432.5
	Davao Orienta					(RCP4.5)	Median Upper Bound	-8.7 20.8	755.5 999.4	-3.3 8.1	591.7 661.4	-5.4 3.6	511.2 560.1	-13.7 -3.8	517.4 576.2
	0 0	827.3	611.8	540.4	599.2		Lower Bound	-17.6	681.4	-10.1	549.9	-12.8	471.2	-23.6	457.9
Н	BVB(High Emission	Median	-2.8	803.8	-1.9	599.9	-5.1	513.0	-6.0	563.0
	7					(RCP8.5)	Upper Bound	18.7	982.0	8.1	661.6	13.4	612.7	6.6	638.5
	~					Moderate	Lower Bound	-13.9	202.7	-12.8	308.1	-15.6	483.0	-28.5	347.6
	North Cotabato					Emission	Median	-1.3	232.4	-8.6	322.8	-5.1	543.1	-15.5	410.4
	Cota	235.4	353.2	572.5	486.0	(RCP4.5)	Upper Bound	7.7	253.4	16.0		11.0	635.2	-4.2	465.5
	£		1,013			High	Lower Bound	-15.9	197.9	-3.0	342.7	-7.7	528.4	-18.7	395.2
112	No					(RCP8.5)	Median	-0.8	233.4	1.2	357.4	1.4	580.5	-7.5	449.3 522.5
Region							Upper Bound Lower Bound	16.9 -24.9	275.2 159.5	19.3 -13.0	421.3 184.9	17.8 -22.6	674.4 258.2	7.5 -31.3	207.7
Re			1			Moderate Emission	Median	-0.3	211.7	-4.5	203.1	1.5	338.7	-10.6	270.4
ĥ	gan	242.0	242.4	222.5	202.5	(RCP4.5)	Upper Bound	19.8	254.3	15.1	244.7	29.0	430.2	14.1	345.0
	Sarangani	212.3	212.6	333.6	302.5	High	Lower Bound	-9.4	192.3	-17.5	175.3	-27.3	242.4	-26.8	221.6
	rv.					Emission	Median	-2.9	206.2	-1.1	210.2	0.9	336.5	-3.0	293.5
	4.5	10000				(RCP8.5)	Upper Bound	12.6	239.1	6.7	226.8	29.6	432.4	5.6	319.4

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-1. CLIRAM of the projected seasonal change in total rainfall (in millimeters) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

	Ĭ.	Obs	erved (1	1971-20	00)				Pro	jected (20	36-2065)				
_	8				-			DJF (Dec	-Jan-Feb)	MAM (Mar	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	MAM	JJA	SON	Scenario	Range*	Percent change	Projected value						
-						Moderate	Lower Bound	-27.3	133.2	-12.9	204.0	-21.6	315.9	-41.4	206.2
	Cotabato					Emission	Median	-8.0	168.7	-7.8	215.8	3.3	416.2	-9.3	318.9
	otat	+00.0	224.4	402.0	251.7	(RCP4.5)	Upper Bound	14.6	210.0	26.1	295.2	40.0	564.0	4.7	368.3
	9	183.3	234.1	402.8	351.7	High	Lower Bound	-20.7	145.3	-9.7	211.3	-16.1	337.9	-23.2	270.1
2	South					Emission	Median	-2.7	178.3	0.1	234.3	1.2	407.6	-3.8	338.4
Region 12	U					(RCP8.5)	Upper Bound	13.9	208.8	6.5	249.4	44.2	580.7	7.6	378.5
egic						Moderate	Lower Bound	-24.0	143.8	-19.2	251.2	-15.7	432.5	-33.8	297.2
æ	Sultan Kudarat					Emission	Median	-10.2	170.0	2.7	319.5	1.1	518.7	-8.6	410.0
	my	+00.0	244.6		****	(RCP4.5)	Upper Bound	10.7	209.6	19.5	371.6	21.6	624.0	3.3	463.5
	- Le	189.3	311.0	513.1	448.7	High	Lower Bound	-22.3	147.1	-8.2	285.5	-17.6	422.9	-21.7	351.3
	Sult					Emission	Median	-5.2	179.5	2.6	319.0	1.9	522.6	-3.6	432.6
Ш	0.					(RCP8.5)	Upper Bound	12.7	213.3	13.8	353.8	36.1	698.5	5.8	474.9
	-ai					Moderate	Lower Bound	-20.3	698.0	-25.3	330.2	-28.6	328.4	-30.4	437.8
	Norte					Emission	Median	-9.8	790.2	-14.2	379.4	-11.4	407.3	-21.2	495.5
	10	025.7	****	460.0	620.0	(RCP4.5)	Upper Bound	31.4	1,150.8	8.3	478.6	1.9	468.7	-1.3	620.6
	Agusan del	875.7	441.9	460.0	628.9	High	Lower Bound	-27.0	639.3	-27.5	320.2	-15.8	387.5	-30.7	435.5
	asní					Emission	Median	0.5	880.2	-9.8	398.7	-10.3	412.8	-10.9	560.5
	A					(RCP8.5)	Upper Bound	20.9	1,058.4	-2.3	431.7	17.6	540.8	8.1	679.6
ľ	Ty 1					Madazita	Lower Bound	-17.7	793.0	-15.3	496.7	-22.2	461.7	-29.7	488.3
	Sur					Moderate Emission	Median	-13.2	835.7	-3.9	563.8	-12.5	518.9	-18.0	569.5
	le le	502.5	233.5	222.0	31.5	(RCP4.5)	Upper Bound	21.4	1,169.0	4.6	613.5	7.1	635.7	-3.2	672.5
	Agusan del	963.3	586.4	593.4	694.8	inc.	Lower Bound	-18.8	782.6	-12.2	514.9	-15.2	503.4	-24.6	524.1
3	gus			-		High Emission	Median	-2.6	937.9	-6.2	550.2	-3.6	572.0	-10.8	619.8
BA	4					(RCP8.5)	Upper Bound	26.3	1,216.8	8.7	637.2	8.4	643.0	9.9	763.5
CARAGA	-61					in a sec	Lower Bound	-12.2	1,239.6	-6.2	600.2	-11.3	397.6	-19.0	678.6
Ü	Norte					Moderate Emission	Median	0.6	1,420.3	3.6	662.7	-4.5	427.6	-14.1	719.6
	del N	J. William	2000	100		(RCP4.5)	Upper Bound	34.9	1,904.3	9.3	699.4	7.1	479.7	8.1	905.4
	0 0	1,412.0	639.6	448.0	837.3	11001	Lower Bound	-19.7	1,134.4	-6.2	599.7	-15.5	378.4	-22.6	648.4
	Surigao		100			High Emission	Median	11.1	1,569.2	-2.8	621.7	0.7	451.0	-11.2	743.6
	Su					(RCP8.5)	Upper Bound	38.9	1,960.6	10.8	709.0	13.1	506.8	15.1	963.5
ľ		-				4000000	Lower Bound	-10.6	1,246.9	-12.9	650.4	-18.5	435.7	-23.2	647.4
	Sur					Moderate Emission	Median	-6.7	1,300.8	-1.7	734.4	-10.6	477.8	-16.9	700.1
		12000	0.30	1000		(RCP4.5)	Upper Bound	27.8	1,781.9	5.6	789.1	-0.2	533.4	-1.0	833.9
	Surigao del	1,394.0	746.9	534.6	842.5		Lower Bound	-15.8	1,173.2	-8.2	685.3	-16.3	447.6	-27.8	608.5
	urlg					High Emission	Median	4.7	1,460.2	-1.8	733.4	-6.4	500.1	-12.0	741.4
	S					(RCP8.5)	Upper Bound	27.5	1,776.9	9.8	820.1	9.6	585.7	8.7	915.9
						in the same of	Lower Bound	-22.0	229.2	-26.1	272.9	-15.0	562.4	-38.8	344.1
	Sur					Moderate Emission	Median	-3.1	284.6	-11.2	327.9	-8.9	602.5	-19.0	455.2
		10000	California of	COL A		(RCP4.5)	Upper Bound	12.3	329.8	13.2	418.2	8.7	719.0	-4.5	537.0
	Lanao del	293.8	369.4	661.5	562.2		Lower Bound	-15.1	249.4	-6.3	346.1	-12.5	579.0	-20.3	447.8
	ana					High Emission	Median	-5.7	277.1	-0.4	368.0	-0.6	657.8	-9.0	511.8
Σ						(RCP8.5)	Upper Bound	12.5	330.6	25.4	463.4	17.3	775.8	7.4	603.8
ARMM						W 100 00 00	Lower Bound	-19.6	181.2	-13.1	346.7	-12.0	558.9	-22.3	430.2
٩	0					Moderate Emission	Median	-3.1	218.4	-4.4	381.4	-1.7	624.8	-10.6	494.7
	lang	1000	1	4.54		(RCP4.5)	Upper Bound	6.9	240.8	22.5	488.8	14.3	726.1	3.9	575.1
	nino	225.3	399.1	635.3	553.6	-	Lower Bound	-19.8	180.6	-0.3	398.1	-12.5	555.9	-15.9	465.3
	Maguindanao	1				High	Median	-4.1	216.1	5.7	421.9	3.1	654.8	-13.9	545.6
	2					(RCP8.5)	-	13.9	256.6	29.6	517.3	8.4	688.9	5.9	586.2
		har at					Upper Bound				317,3	0.4	000.9	5.9	300.2

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperaturel (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000.

Ŧ		Obs	served (1	1971-20	00)	7			Pro	jected (20	036-2065)	-			
_	8							DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	MAM	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
1						Moderate	Lower Bound	0.9	26.2	0.9	29.0	1.0	29,3	1.0	28.4
	ie i					Emission	Median	1.2	26.5	1.2	29.3	1.3	29.6	1.1	28.5
	No	25.3	28.1	28.3	27.4	(RCP4.5)	Upper Bound	1.6	26.9	1.7	29.8	1,8	30.1	1.8	29.2
	Ilocos Norte	25.5	20.1	20.5	27.17	High	Lower Bound	1.1	26.4	1.3	29.4	1.3	29.6	1.3	28.7
	oll lie					Emission	Median	1.6	26.9	1.6	29.7	1.6	29.9	1.5	28.9
į,						(RCP8.5)	Upper Bound	1.9	27.2	2.2	30.3	2.4	30.7	2.2	29.6
	111					Moderate	Lower Bound	1.1	24.2	1.0	26.7	1.0	26.4	1.0	25.8
	Sur					(RCP4.5)	Median	1.3	24.4	1.3	27.0	1.3	26.7	1.1	25.9
	So	23.1	25.7	25.4	24.8	(MCI 415)	Upper Bound	1.9	25.0	1.8	27.5	1.8	27.2	1.9	26.7
	llocos!					High	Lower Bound	1.2	24.3	1.4	27.1	1.3	26.7	1.4	26.2
11						(RCP8.5)	Median Upper Bound	2.1	24.8	1.6	27.3	2.3	27.0	2.4	26.4
Region							Lower Bound	1.1	21.6	1.0	23.9	0.9	23.7	1.0	23.2
Re						Moderate Emission	Median	1.3	21.8	1.2	24.1	1.2	24.0	1.0	23.2
	lon	1000	1000	4.5		(RCP4.5)	Upper Bound	1.8	22.3	1.7	24.6	1.8	24.6	1.8	24.0
	La Union	20.5	22.9	22.8	22.2		Lower Bound	1.2	21.7	1.3	24.2	1.3	24.1	1.3	23.5
	3					High Emission	Median	1.6	22.1	1.6	24.5	1.5	24.3	1.6	23.8
						(RCP8.5)	Upper Bound	2.1	22.6	2.3	25.2	2.3	25.1	2.3	24.5
1						Madanika	Lower Bound	1.1	26.1	1.0	28.4	1.0	27.9	1.0	27.4
	ь					Moderate Emission	Median	1.3	26.3	1.2	28.6	1.3	28.2	1.1	27.5
	Pangasinan	75.0	22.4	20.0	20.4	(RCP4,5)	Upper Bound	1.8	26.8	1.8	29.2	1.9	28.8	1.9	28.3
	nga	25.0	27.4	26.9	26.4	High	Lower Bound	1.3	26.3	1.4	28.8	1.3	28.2	1.4	27.8
	Pa					Emission	Median	1.6	26.6	1.7	29.1	1.6	28.5	1.6	28.0
						(RCP8,5)	Upper Bound	2.1	27.1	2.4	29.8	2.4	29.3	2.3	28.7
						Moderate	Lower Bound	1.0	25.5	1.0	28.4	1.0	28.2	1.0	27.4
			10.11			Emission	Median	1.2	25.7	1.2	28.6	1.3	28.5	1.2	27.6
	Abra	24.5	27.4	27.2	26.4	(RCP4,5)	Upper Bound	1.7	26.2	1.8	29.2	1.8	29.0	1.8	28.2
	¥	24.5	27.5	27.2	20.4	High	Lower Bound	1.2	25.7	1.4	28.8	1.4	28.6	1.4	27.8
						Emission	Median	1.7	26.2	1.7	29.1	1.6	28.8	1.7	28.1
100	,					(RCP8.5)	Upper Bound	2.0	26.5	2,3	29.7	2.4	29.6	2.3	28.7
						Moderate	Lower Bound	0.9	25.7	0.8	28.8	1.0	29.4	0.9	28.0
						(RCP4.5)	Median	1.2	26.0	1.2	29.2	1.2	29.6	1,1	28.2
	Apayao	24.8	28.0	28.4	27.1	(RCF4.3)	Upper Bound	1.4	26.2	1,6	29.6	1.8	30.2	1.8	28.9
	Ap	1				High	Lower Bound	1.0	25.8	1,2	29.2	1,4	29.8	1,3	28.4
						(RCP8.5)	Median	1.5	26.3	1.7	29.7	2.4	30.0	1.5	28.6
1						1000	Upper Bound Lower Bound	1.7	26.5	2.3	30.3 23.0	1.1	30.8	2.1	29.2
						Moderate Emission	Median Median	1.3	20.7	1.3	23.2	1.4	23.4	1.3	22.5
	iğ	14.00	2.50	22.7		(RCP4.5)	Upper Bound	1.8	21.2	1.9	23.8	1.9	23.9	1.9	23.1
	Benguet	19.4	21.9	22.0	21.2	10000	Lower Bound	1.3	20.7	1.5	23.4	1.4	23.4	1.4	22.6
	m					High Emission	Median	1.8	21.2	1.8	23.7	1.7	23.7	1.7	22.9
8						(RCP8.5)	Upper Bound	2.1	21.5	2.4	24.3	2.5	24.5	2.3	23.5
CAR						Moderate	Lower Bound	1.0	23.2	0.9	26.5	1.0	26.8	1.0	25.5
						Emission	Median	1.2	23.4	1.2	26.8	1.3	27,1	1.1	25.6
	390	22.7	25.6	25.0	74.5	(RCP4.5)	Upper Bound	1.5	23.7	1.7	27.3	1.9	27.7	1.9	26.4
	Ifugao	22.2	25.6	25.8	24.5	High	Lower Bound	1.1	23.3	1.3	26.9	1.4	27,2	1.3	25.8
						Emission	Median	1.6	23.8	1.7	27.3	1.6	27.4	1.6	26.1
	Щ					(RCP8.5)	Upper Bound	1.8	24.0	2.3	27.9	2.5	28.3	2.3	26.8
	7.11		-			Moderate	Lower Bound	1.0	24.8	0.9	28.4	1.0	28.7	1.0	27.1
			100			Emission	Median	1.2	25.0	1.2	28.7	1,3	29.0	1.1	27.2
	Kalinga	23.8	27.5	27.7	26.1	(RCP4.5)	Upper Bound	1.5	25.3	1.7	29.2	1.9	29.6	1.8	27.9
	Kal	23,0	27,3	21.10	20.1	High	Lower Bound	1.1	24.9	1.2	28.7	1.4	29.1	1.3	27.4
						Emission	Median	1.6	25.4	1.7	29.2	1.6	29.3	1.6	27.7
	2.1	11-11				(RCP8.5)	Upper Bound	1.7	25.5	2.3	29.8	2.4	30.1	2.2	28.3
	a)Ce					Moderate	Lower Bound	1.0	23.7	1.0	27.0	1.0	27.1	1.0	25.9
	ovin		100			Emission (RCP4.5)	Median	1.2	23.9	1.2	27.2	1.3	27.4	1,2	26.1
	P.	22.7	26.0	26.1	24.9	(KCP4.5)	Upper Bound	1.6	24.3	1.8	27.8	2.0	28.1	1.9	26.8
	Mountain Province		75.75)	7343	0.102	High	Lower Bound	1.2	23.9	1.3	27.3	1.3	27.4	1.4	26.3
	loui					(RCP8.5)	Median	1,7	24.4	1.7	27.7	1.6	27.7	1.6	26.5
	20					(ACLOS)	Upper Bound	1.9	24.6	2.3	28.3	2.5	28.6	2.3	27.2

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperature (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

		Obs	served (1	971-200	00)	11			Pro	jected (2	036-2065)				
_	aor l	1				11		DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	МАМ	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
ľ						Moderate	Lower Bound	0.9	25.4	0.9	29.0	1.0	29.9	0.9	28.0
	e e					(RCP4.5)	Median	1.2	25.7	1.2	29.3	1.3	30.2	1.1	28.2
	Cagayan	24.5	28.1	28.9	27.1	(rici ris)	Upper Bound	1.4	25.9 25.5	1.7	29.8	1.9	30.8	1.8	28.9
	8	I Kar		1.0		High Emission	Lower Bound Median	1.5	26.0	1.6	29.3	1.6	30.5	1.5	28.6
						(RCP8.5)	Upper Bound	1.7	26.2	2.2	30.3	2.4	31.3	2.2	29.3
-		-				Moderate	Lower Bound	1.0	25.1	0.9	28.8	1.0	29.7	1.0	27.8
						Emission	Median	1.2	25.3	1.2	29.1	1.3	30.0	1.1	27.9
	Isabela	24.1	27.9	28.7	26.8	(RCP4.5)	Upper Bound	1.5	25.6	1.7	29.6	2.0	30.7	1.9	28.7
	Isa		41.10	1000	-	High	Lower Bound	1.1	25.2	1.2	29.1	1.3	30.0	1.3	28.1
1 2						(RCP8.5)	Median	1.6	25.7	1.7	29.6	1.6	30.3	1.6	28.4
Region	-			-			Upper Bound Lower Bound	1.8	25.9 23.3	2.3	30.2 26.1	2.5	31.2 26.4	2.3	29.1 25.4
Re	Pλ					Moderate Emission	Median.	1.2	23.5	1.3	26.4	1.3	26.7	1.1	25.5
	Nueva Vizcaya	1200	2507	200	40.00	(RCP4.5)	Upper Bound	1.6	23.9	1.8	26.9	1.9	27.3	1.9	26.3
	va v	22.3	25.1	25.4	24.4	High	Lower Bound	1.1	23.4	1.3	26.4	1.4	26.8	1.4	25.8
	Nue		1.0			Emission	Median	1.6	23.9	1.7	26.8	1.6	27,0	1.6	26.0
						(RCP8.5)	Upper Bound	1.9	24.2	2.3	27.4	2.5	27.9	2.3	26.7
						Moderate	Lower Bound	1.0	24.7	1.0	27.8	1.0	28.6	1.0	27.2
	0			100		(RCP4.5)	Median	1.2	24.9	1.3	28.1	1.3	28.9	1.1	27.3
	Quirino	23.7	26.8	27.6	26.2	State stary	Lower Bound	1.6	25.3 24.8	1.8	28.6 28.1	2.0	29.6	1.9	28.1
	o.	1		-		High Emission	Median	1.5	25.2	1.7	28.5	1.6	29.0	1.6	27.8
						(RCP8,5)	Upper Bound	1.9	25.6	2.3	29.1	2.5	30.1	2.3	28.5
						Moderate	Lower Bound	1.0	25.5	0.9	28.0	1.0	28.9	1.0	27.7
	1					Emission	Median	1.2	25.7	1.2	28.3	1.2	29.1	1.1	27.8
	Aurora	24.5	27.1	27.9	26.7	(RCP4.5)	Upper Bound	1.6	26.1	1.7	28.8	1.9	29.8	1.9	28.6
	Au	2113	-7.12	2012	2017	High	Lower Bound	1.1	25.6	1.3	28.4	1.4	29.3	1.3	28.0
						(RCP8.5)	Median	1.6	26.1	1.6	28.7	1.6	29.5	1.5	28.2
b				-		(merono)	Upper Bound	1.9	26.4 27.4	0.9	29.2	1.0	30.3	1.0	28.9
						Moderate Emission	Lower Bound Median	1.2	27.4	1.2	29.6	1.2	28.6	1.0	28.3
	up up	600	23.2	4		(RCP4.5)	Upper Bound	1.7	28.1	1.8	30.5	1.7	29.3	1.8	29.1
	Bataan	26.4	28.7	27.6	27.3	mak	Lower Bound	1.2	27.6	1.3	30.0	1.3	28.9	1.4	28.7
	-					High Emission	Median	1.6	28.0	1.7	30.4	1.5	29.1	1.5	28.8
						(RCP8.5)	Upper Bound	2.0	28.4	2.3	31.0	2.2	29.8	2.2	29.5
						Moderate	Lower Bound	1.0	26.6	1.0	28.9	1.0	28.1	1.0	27.7
	-					(RCP4.5)	Median	1.2	26.8	1,2	29.1	1.3	28.4	1,1	27.8
	Bulacan	25.6	27.9	27.1	26.7	(1101-110)	Upper Bound	1.6	27.2	1.7	29.6	1.8	28.9	1.9	28.6
	B	1111		***		High Emission	Lower Bound Median	1.2	26.8	1.7	29.2	1.6	28.4	1.4	28.1
ļ,						(RCP8.5)	Upper Bound	1.9	27.5	2.3	30.2	2.3	29.4	2.3	29.0
b	-	-				Moderate	Lower Bound	1.0	26.3	1.0	28.7	1.0	28.5	1.0	27.8
3	ja ja					Emission	Median	1.2	26.5	1,2	28.9	1.3	28.8	1,1	27.9
uo	Eci	25.3	27.7	27.5	26.8	(RCP4.5)	Upper Bound	1.7	27.0	1.8	29.5	1.9	29.4	1.9	28.7
Region	Nueva Ecija	25.5	27.17	27.5	20.0	High	Lower Bound	1.2	26.5	1.4	29,1	1.3	28.8	1,4	28.2
	Z					Emission (RCP8.5)	Median	1.6	26.9	1.7	29.4	1.6	29.1	1.6	28.4
ļ,						(10.00)	Upper Bound	2.0	27.3	2.3	30.0	2.4	29.9	2.3	29,1
	4					Moderate Emission	Lower Bound Median	1.0	27.0	1.0	29.3	1.0	28.5	1.0	28.1
	ınga	75.00	22.2	777		(RCP4.5)	Upper Bound	1.7	27.7	1.8	30.1	1.9	29.4	1.9	29.0
	Pampanga	26.0	28.3	27.5	27.1	Disc	Lower Bound	1.2	27.2	1.3	29.6	1.4	28.9	1.4	28.5
	Pa					High Emission	Median	1.6	27.6	1.7	30.0	1.6	29.1	1.6	28.7
	20					(RCP8.5)	Upper Bound	2.1	28.1	2.4	30.7	2.4	29.9	2.3	29.4
	1					Moderate	Lower Bound	1.0	27.1	1.0	29.3	1.0	28.8	1.0	28.3
						Emission (RCP4.5)	Median	1.3	27.4	1.2	29.5	1.2	29.0	1.1	28.4
	Tarlac	26.1	28.3	27.8	27.3	(ACP4.3)	Upper Bound	1.7	27.8	1.8	30.1	1.9	29.7	1.9	29.2
	P	127				High	Lower Bound	1.2	27.3	1.4	29.7	1.4	29.2	1.4	28.7
						(RCP8.5)	Median Upper Bound	1.6	27.7	2.3	30.0 30.6	1.6 2.5	29.4 30.3	1.6 2.3	28.9
							Lower Bound	1.0	27.3	1.0	29.3	1.0	28.4	1.0	28.2
	4					Moderate Emission	Median	1.3	27.6	1.2	29.5	1.2	28.6	1.1	28.3
	ales	76.7	20.2	27.4	27.0	(RCP4.5)	Upper Bound	1.7	28.0	1.8	30.1	1.8	29.2	1.8	29.0
	Zambales	26.3	28.3	27,4	27.2	High	Lower Bound	1.2	27.5	1.3	29.6	1.4	28.8	1.4	28.6
	И					Emission	Median	1.6	27.9	1.7	30.0	1.5	28.9	1.6	28.8
Н	4.13					(RCP8.5)	Upper Bound	2.1	28.4	2.3	30.6	2.2	29.6	2.2	29.4

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperature (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

Ī		Obs	served (1	1971-20	00)	71	,				36-2065)				
Ē	nce	4:5		7.4		3	2000	DJF (Dec	-Jan-Feb)	MAM (Mar	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	МАМ	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
T						Moderate	Lower Bound	1.0	25.2	1.0	27.5	1.0	26.9	1.0	26.6
	se		100			(RCP4.5)	Median	1.2	25.4	1.2	27.7	1.2	27.1	1.0	26.6
	Batangas	24.2	26.5	25.9	25.6	y said they	Upper Bound Lower Bound	1.6	25.8 25.4	1.7	28.2	1.7	27.6 27.2	1.8	27.4
	Ba		TO Y			High Emission	Median	1.6	25.8	1.6	28.1	1.5	27.4	1.5	27.1
	15					(RCP8.5)	Upper Bound	1.9	26.1	2.2	28.7	2.3	28.2	2.2	27.8
						Moderate	Lower Bound	1.0	26.7	1.0	29.2	1.0	28.3	1.0	27.9
						Emission	Median	1.3	27.0	1.2	29.4	1.2	28.5	1.1	28.0
	Cavite	25.7	28.2	27.3	26.9	(RCP4.5)	Upper Bound	1.7	27.4	1.7	29.9	1.8	29.1	1.8	28.7
	S	3446	120.01	AACC4		High	Lower Bound	1.2	26.9	1.3	29.5	1.3	28.6	1.4	28.3
						(RCP8.5)	Median Upper Bound	2.0	27.3	2.3	29.8 30.5	2.2	28.8	1.5 2.2	28.4
P				-		400000000000000000000000000000000000000	Lower Bound	1.0	26.0	1.0	28.5	1.0	28.5	1.0	27.7
4						Moderate Emission	Median	1.2	26.2	1.2	28.7	1.3	28.8	1.1	27.8
4	Laguna	25.0	27.5	27.5	26.7	(RCP4.5)	Upper Bound	1.6	26.6	1.7	29.2	1.8	29.3	1.9	28.6
Region	Lag	25.0	27,3	27.5	20.7	High	Lower Bound	1.1	26.1	1.3	28.8	1.3	28.8	1.4	28.1
ď						Emission (RCP8.5)	Median	1.6	26.6	1.6	29.1	1.5	29.0	1.5	28.2
ŝ						(KCF6.5)	Upper Bound	1.9	26.9	2.2	29.7	2.4	29.9	2.3	29.0
			-			Moderate	Lower Bound	1.0	26.1	0.9	28.1	1.0	28.6	1.0	27.7
	6		22.2	dea.		(RCP4,5)	Median Upper Bound	1.6	26.7	1.6	28.8	1.8	29.4	1.8	28.5
	Quezor	25.1	27.2	27.6	26.7	were .	Lower Bound	1.1	26.2	1.3	28.5	1.3	28.9	1.3	28.0
						High Emission	Median	1.6	26.7	1.6	28.8	1.5	29.1	1.4	28.1
						(RCP8.5)	Upper Bound	1.8	26.9	2.0	29.2	2.3	29.9	2.2	28.9
						Moderate	Lower Bound	1.0	26.4	1.0	28.9	1.0	28.6	1.0	27.8
	12					(RCP4.5)	Median	1.2	26.6	1.2	29.1	1.3	28.9	1.1	27.9
	Rizal	25.4	27.9	27.6	26.8	(1501-4.5)	Upper Bound	1.6	27.0	1.7	29.6	1.8	29.4	1.8	28.6
		200	100	***		High Emission	Lower Bound Median	1.2	26.6	1.6	29.2	1.5	28.9	1.4	28.2
						(RCP8.5)	Upper Bound	1.9	27.3	2.2	30.1	2.3	29.9	2.2	29.0
	o	1				Madausta	Lower Bound	1.0	27.5	1.0	29.3	0.9	28.2	1.0	28.1
	ndor					Moderate Emission	Median	1.3	27.8	1.3	29.6	1.2	28.5	1.1	28.2
	Occidental Mindoro	26.5	28.3	27.3	27.1	(RCP4.5)	Upper Bound	1.7	28.2	1.8	30.1	1.7	29.0	1.8	28.9
	ents	20.5	20.5	27.5	27.1	High	Lower Bound	1.3	27.8	1.4	29.7	1.4	28.7	1.4	28.5
	ccid					(RCP8.5)	Median	1.6	28.1	1.7	30.0	1.5	28.8	1.5	28.6
p	0					(Mar als)	Upper Bound	1.0	28.5 27.4	1.0	30.6 29.3	0.9	29.5	1.0	29.3
	oro					Moderate Emission	Lower Bound Median	1.2	27.4	1.2	29.5	1.2	28.8	1.1	28.4
	Oriental Mindoro	7.4		-328	22.2	(RCP4.5)	Upper Bound	1.6	28.0	1.7	30.0	1.8	29.4	1.8	29.1
	tal	26.4	28.3	27.6	27.3	High	Lower Bound	1.2	27.6	1,3	29.6	1.3	28.9	1.3	28.6
4-B	rien					Emission	Median	1.6	28.0	1.6	29.9	1.5	29.1	1.5	28,8
on 4	Ü					(RCP8.5)	Upper Bound	1.9	28.3	2,2	30.5	2.3	29.9	2,2	29.5
Region						Moderate	Lower Bound	1.0	27.9	1.0	29.1	0.9	28.2	1.0	28.4
•	ug.			100		(RCP4.5)	Median	1.3	28.2	1.2	29.3	1.2	28.5	1.1	28.5
	Palawan	26.9	28.1	27.3	27.4		Upper Bound Lower Bound	1.6	28.5	1.6	29.7	1.6	28.9 28.6	1.7	29.1
	Pa					High Emission	Median	1.6	28.5	1.7	29.8	1.5	28.8	1.5	28.9
						(RCP8.5)	Upper Bound	1.9	28.8	2.1	30.2	2.0	29.3	2.1	29.5
	111					Moderate	Lower Bound	0.9	27.2	1.0	29.5	0.9	29.0	1.0	28.7
	5					Emission	Median	1.2	27.5	1.2	29.7	1.2	29.3	1.1	28.8
	Romblon	26.3	28.5	28.1	27.7	(RCP4.5)	Upper Bound	1.5	27.8	1.5	30.0	1.7	29.8	1.7	29.4
	Ron	10000		21.5		High	Lower Bound	1.1	27.4	1.3	29.8	1.3	29.4	1.3	29.0
						(RCP8.5)	Median	1.5	27.8 28.0	1.6	30.1	1.4	29.5 30.2	2.1	29.1 29.8
-	-						Upper Bound Lower Bound	1.0	27.1	0.9	29.7	1.0	29.0	1.0	29.8
	in					Moderate Emission	Median	1.2	27.3	1.2	30.0	1.3	29.3	1.1	28.5
2	Metro Manila	20.4	20.0	20.0	22.0	(RCP4.5)	Upper Bound	1.6	27.7	1.7	30.5	1.8	29.8	1.8	29.2
NCR	tro	26.1	28.8	28.0	27.4	High	Lower Bound	1.2	27.3	1.3	30.1	1.3	29.3	1.3	28.7
	M					Emission	Median	1.6	27.7	1.6	30.4	1.5	29.5	1.5	28.9
						(RCP8.5)	Upper Bound	1.9	28.0	2.2	31.0	2.3	30.3	2.2	29.6
						Moderate	Lower Bound	1.0	26.6	1.0	28.2	1.0	28.8	1.0	28.1
2			2.3			(RCP4.5)	Median	1.2	26.8	1.2	28.4	1.2	29.0	1.1	28.2
Region	Albay	25.6	27.2	27.8	27.1		Lower Bound	1.7	27.3 26.8	1.8	29.0 28.6	1.8	29.6 29.2	1.8	28.9 28.5
Re	-					High Emission	Median	1.6	27.2	1.7	28.9	1.6	29.4	1.5	28.6
						(RCP8.5)	Upper Bound	2.1	27.7	2.3	29.5	2.2	30.0	2.2	29.3

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperature (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

Ħ		Obs	erved (1	971-200	00)	21			Pro	jected (2	36-2065)				
	8	1		100				DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	MAM	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
	81					Moderate	Lower Bound	1.0	26.7	1.0	28.6	1.0	29.3	1.0	28.3
	Camarines Norte	1	lane.			Emission (RCP4.5)	Median	1.2	26.9	1.2	28.8	1.2	29.5	1.0	28.3
	nes	25.7	27.6	28.3	27.3	(RCP4.5)	Upper Bound	1.6	27.3	1.7	29.3	1.8	30,1	1.8	29.1
	mari	100	-	-		High	Lower Bound	1.2	26.9	1.3	28.9	1.3	29.6	1.3	28.6
	Š			-		(RCP8.5)	Median Upper Bound	1.9	27.6	2.2	29.2	2.3	30.6	2.2	29.5
-							Lower Bound	1.0	26.6	1.0	28.5	1.0	29.1	1.0	28.1
	Sur					Moderate Emission	Median	1.2	26.8	1.2	28.7	1.2	29.3	1.1	28.2
		25.6	27.5	28.1	27.1	(RCP4.5)	Upper Bound	1.6	27.2	1.7	29.2	1.8	29.9	1.8	28.9
	Camarines	25.0	27.5	20.1	27,1	High	Lower Bound	1.2	26.8	1.3	28.8	1.4	29.5	1.4	28.5
	3					Emission (RCP8.5)	Median	1.6	27.2	1.6	29.1	1.5	29.6	1.5	28.6
1,						(KCF8,5)	Upper Bound	1.9	27.5	2.2	29.7	2.2	30.3	2.2	29.3
	- UT					Moderate	Lower Bound Median	1.0	25.7 25.8	1.0	27.3 27.5	1.0	28.2	1.0	27.4 27.5
n 5	Catanduanes		1,000	2.0		(RCP4.5)	Upper Bound	1.5	26.2	1.6	27.9	1.7	28.9	1.7	28.1
Region	andı	24.7	26.3	27.2	26.4	COSA	Lower Bound	1.2	25.9	1.3	27.6	1.3	28.5	1.3	27.7
æ	Cata					High Emission	Median	1.5	26.2	1.6	27.9	1.6	28.8	1.4	27.8
ľ						(RCP8.5)	Upper Bound	1.7	26.4	1.9	28.2	2.1	29.3	2.0	28.4
						Moderate	Lower Bound	1.0	27.6	1.0	29.4	1.0	29.6	1.0	29.0
	-0					Emission (RCP4,5)	Median	1.2	27.8	1.2	29.6	1.2	29.8	1.1	29.1
	Masbate	26.6	28.4	28.6	28.0	(KCP4,5)	Upper Bound	1.6	28.2	1.6	30.0	1.7	30.3	1.7	29.7
	Ma	00.0	1000	-		High	Lower Bound	1.2	27.8	1.3	29.7 30.0	1.3	29.9	1.3	29.3
						(RCP8,5)	Median Upper Bound	1.8	28.4	1.9	30.3	2.1	30.1	2.1	30.1
r							Lower Bound	1.0	26.9	1.0	28.4	1.0	28.9	1.0	28.3
	_					Moderate Emission	Median	1.2	27.1	1.2	28.6	1.2	29.1	1.1	28.4
	Sorsogon	25.0	27.4	27.0	27.2	(RCP4.5)	Upper Bound	1.6	27.5	1.6	29.0	1.7	29.6	1.8	29.1
	Sors	25.9	27.4	27.9	27.3	High	Lower Bound	1.2	27.1	1.4	28.8	1.4	29.3	1.3	28.6
	0,					Emission	Median	1.5	27.4	1.6	29.0	1.6	29.5	1.5	28.8
÷						(RCP8.5)	Upper Bound	1.9	27.8	2.0	29.4	2.2	30.1	2.2	29.5
Ī						Moderate	Lower Bound	1.0	27.1	1.0	28.9	0.9	28.7	1.0	28.4
	-		0.7	100		(RCP4.5)	Median	1.2	27.3	1.2	29.1	1.1	28.9	1.1	28.5
	Aklan	26.1	27.9	27.8	27.4	AC 15-2	Upper Bound Lower Bound	1.2	27.7	1.3	29.3	1.3	29.0	1.3	28.7
						High Emission	Median	1.5	27.6	1.6	29.5	1.5	29.3	1.4	28.8
						(RCP8.5)	Upper Bound	1.8	27.9	2.1	30.0	2.2	30.0	2.2	29.6
						Moderate	Lower Bound	1.0	27.6	1.0	29.4	0.9	28.8	1.0	28.7
						Emission	Median	1.3	27.9	1.3	29.7	1.2	29.1	1,1	28.8
	Antique	26.6	28.4	27.9	27.7	(RCP4.5)	Upper Bound	1.7	28.3	1.8	30.2	1.8	29.7	1.8	29.5
	An	0000	1,000.00	232		High	Lower Bound	1.2	27.8	1.4	29.8	1,4	29.3	1.4	29.1
			-			(RCP8.5)	Median Upper Bound	2.0	28.2	1.8	30.2	1.5 2.3	29.4 30.2	1.5	29.2
Ŕ						Total State of the last	Lower Bound	1.0	26.9	1.0	28.7	1.0	28.8	1.0	28.3
						Moderate Emission	Median	1.2	27.1	1,2	28.9	1.2	29.0	1.1	28.4
	zic		200	-56	000	(RCP4.5)	Upper Bound	1.6	27.5	1.7	29.4	1.9	29.7	1.8	29.1
	Capiz	25.9	27.7	27.8	27.3	High	Lower Bound	1.2	27.1	1.4	29.1	1,4	29.2	1.3	28.6
9						Emission	Median	1.5	27.4	1.6	29.3	1.6	29.4	1,5	28.8
Region						(RCP8.S)	Upper Bound	1.9	27.8	2.2	29.9	2.3	30.1	2,2	29.5
Reg						Moderate	Lower Bound	1.0	27.5	1.0	29.3	0.9	28.6	1.0	28.4
Ĩ	ras					(RCP4.5)	Median	1.2	27.7 28.0	1.2	29.5	1.1	28.8	1.1	28.5
	Guimaras	26.5	28.3	27.7	27.5		Upper Bound Lower Bound	1.5	27.8	1.5	29.8 29.6	1.3	29.2	1.6	29.1
	5			777		High Emission	Median	1.5	28.1	1.6	29.9	1.4	29.1	1.4	28.9
	1					(RCP8.5)	Upper Bound	1.8	28.4	2.1	30.4	1.9	29.6	2.0	29.5
	1					Moderate	Lower Bound	1.0	27.4	1.0	29.2	0.9	28.8	1.0	28.6
						Emission	Median	1.2	27.6	1.2	29.4	1.2	29.1	1.1	28.7
	tiolio	26.4	28.2	27.9	27.6	(RCP4.5)	Upper Bound	1.6	28.0	1.7	29.9	1.8	29.7	1.8	29.4
	II	200	2000	200	21.16	High	Lower Bound	1.2	27.6	1.3	29.5	1.4	29.3	1.3	28.9
						(RCP8.5)	Median	1.6	28.0	1.7	29.9	1.5	29.4	1.5	29.1
						200000	Upper Bound	1.9	28.3	1.0	30.4 29.4	1.0	30.1 28.8	1.0	29.8
	unta					Moderate Emission	Lower Bound Median	1.0	27.9	1.3	29.4	1.0	29.0	1.1	28.7
	cide	Carried Street	1000	1576	73.4	(RCP4.5)	Upper Bound	1.7	28.4	1.8	30.2	1.8	29.6	1.8	29.4
	Negros Occidental	26.7	28.4	27.8	27.6	High	Lower Bound	1.3	28.0	1.3	29.7	1.4	29.2	1.4	29.0
	egro					Emission	Median	1.6	28.3	1.7	30.1	1.6	29.4	1.5	29.1
	ž		-			(RCP8.5)	Upper Bound	2.1	28.8	2.3	30.7	2.3	30.1	2.2	29.8

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperature (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

Ī		Obs	served (1	971-200	00)		>		Pro	jected (20	36-2065)				
_	90	-				Trans.		DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	МАМ	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
Ī						Moderate	Lower Bound	1.0	27.6	1.0	29.0	1,0	29.2	1.0	28.8
	Ш		1000			(RCP4.5)	Median	1.2	27.8	1.2	29.2	1.2	29.4	1.2	29.0
	Bohol	26.6	28.0	28.2	27.8	(mar may	Upper Bound	1.7	28.3	1.7	29.7	1.7	29.9	1.8	29.6 29.2
			100	100		High Emission	Lower Bound Median	1.6	28.2	1.7	29.4	1.6	29.8	1.4	29.2
ı						(RCP8.5)	Upper Bound	2.0	28.6	2.2	30.2	2.2	30.4	2.2	30.0
ř	-					Moderate	Lower Bound	1.0	27.8	1.0	29.4	0.9	29.1	1.0	28.9
1						Emission	Median	1.2	28.0	1.2	29.6	1.2	29.4	1.1	29.0
5	Cebu	26.8	28.4	28.2	27.9	(RCP4.5)	Upper Bound	1.7	28.5	1.7	30.1	1.7	29.9	1.8	29.7
Region	ŭ	20.0	20.4	20,2	21.13	High	Lower Bound	1.3	28.1	1.3	29.7	1.4	29.6	1.3	29.2
7						(RCP8.5)	Median	1.6	28.4	1.6	30.0	1.6	29.8	1.5	29.4
ŀ				-		(NCFO.5)	Upper Bound	2.0	28.8	2.2	30.6	2.2	30.4	2.2	30.1
	豆					Moderate	Lower Bound Median	1.0	28.0	1.0	29.4	1.2	28.9	1.0	28.8
	Negros Oriental		4.0	0.00		(RCP4.5)	Upper Bound	1.6	28.6	1.7	30.1	1.7	29.7	1.7	29.5
	0 80	27.0	28.4	28.0	27.8		Lower Bound	1.3	28.3	1.4	29.8	1.4	29.4	1.3	29.1
	egro		100			High Emission	Median	1.6	28.6	1.7	30.1	1.5	29.5	1.5	29.3
	Z					(RCP8.5)	Upper Bound	2.0	29.0	2.3	30.7	2.2	30.2	2.2	30.0
Ŧ	10	- 1				Moderate	Lower Bound	1.0	27.1	1.0	28.7	1.0	29.3	1.0	28.7
	Samar					Emission	Median	1.1	27.2	1.2	28.9	1.2	29.5	1.2	28.9
Н	Sa	26.1	27.7	28.3	27.7	(RCP4,5)	Upper Bound	1.6	27.7	1.7	29.4	1.8	30.1	1.8	29.5
	Eastern	20.1	27.17	20,5	27.7	High	Lower Bound	1.3	27.4	1.4	29.1	1.4	29.7	1.4	29.1
ľ	Ea					Emission (RCP8,5)	Median	1.5	27.6	1.6	29.3	1.7	30.0	1.5	29.2
,						(ACFO,3)	Upper Bound	1.9	28.0	2.1	29.8	2.2	30.5	2.2	29.9
						Moderate	Lower Bound	1.0	27.4 27.6	1.0	28.8	0.9	28.9	1.0	28.7
	0		100	1000		(RCP4.5)	Median Upper Bound	1.2	28.1	1.2	29.0 29.6	1.2	29.2	1.1	28.8
	Leyte	26.4	27.8	28.0	27.7	and the same of th	Lower Bound	1.3	27.7	1.4	29.2	1.4	29.4	1.4	29.1
	- 1		100			High Emission	Median	1.5	27.9	1.7	29.5	1.6	29.6	1.5	29.2
						(RCP8.5)	Upper Bound	2.1	28.5	2.3	30.1	2.2	30.2	2.3	30.0
ì	7					Moderate	Lower Bound	1.0	27.0	1.0	28.5	1.0	29.3	1.0	28.5
8	Samar					Emission	Median	1.2	27.2	1.2	28.7	1.2	29.5	1.1	28.6
Region	n Sa	26.0	27.5	28.3	27.5	(RCP4.5)	Upper Bound	1.6	27.6	1.7	29.2	1.8	30.1	1.8	29.3
Seg	Northern	20.0	27.3	20.5	27.5	High	Lower Bound	1.3	27.3	1.4	28.9	1.4	29.7	1.4	28.9
_	Nort					Emission	Median	1.5	27.5	1.6	29.1	1.6	29.9	1.5	29.0
j.						(RCP8.5)	Upper Bound	1.9	27.9	2,2	29.7	2.3	30.6	2.2	29.7
١						Moderate	Lower Bound	1.0	27.3	1.0	28.9	1.0	29.4	1.0	28.8
	L .		700			(RCP4.5)	Median	1.2	27.5 28.0	1,2	29.1	1.2	29.6 30.2	1.2	29.0
ı	Samar	26.3	27.9	28.4	27.8		Upper Bound Lower Bound	1.7	27.6	1.4	29.7	1.4	29.8	1.4	29.2
	U.					High Emission	Median	1.5	27.8	1.7	29.6	1.6	30.0	1.6	29.4
ı						(RCP8.5)	Upper Bound	2.1	28.4	2.3	30.2	2.3	30.7	2.3	30.1
ì	-					Moderate	Lower Bound	1.0	27.4	1.0	28.7	0.9	28.7	1.0	28.5
	eyte					Emission	Median	1.2	27.6	1.2	28.9	1.2	29.0	1.1	28.6
	Southern Leyte	26.4	27.7	27.8	27.5	(RCP4.5)	Upper Bound	1.7	28.1	1.7	29.4	1.7	29.5	1.8	29.3
	athe	20.7	20.0	27.0	27.13	High	Lower Bound	1.3	27.7	1.3	29.0	1.3	29.1	1.3	28.8
	Sol					(RCP8.5)	Median	1.5	27.9	1.6	29.3	1.6	29.4	1.5	29.0
						(Upper Bound	2.0	28.4	2.1	29.8	2.1	29.9	2.2	29.7
1	del					Moderate Emission	Lower Bound Median	0.9	27.9 28.2	1.0	28.9	1.2	28.5	1.0	28.5
ľ	egr.		33.7			(RCP4.5)	Upper Bound	1.7	28.2	1.7	29.2	1.8	29.4	1.8	29.3
	Zamboanga Norte	27.0	27.9	27.6	27.5	Thee	Lower Bound	1.4	28.4	1.4	29.3	1.3	28.9	1.3	28.8
ı	Cam					High Emission	Median	1.6	28.6	1.7	29.6	1.5	29.1	1.5	29.0
	2					(RCP8.5)	Upper Bound	2.1	29.1	2.2	30.1	2.3	29.9	2.2	29.7
ľ	71					Moderate	Lower Bound	1.0	27.8	1.1	28.7	1.0	28.3	1.0	28.2
6	Zamboanga del Sur					Emission	Median	1.2	28.0	1.3	28.9	1.2	28.5	1.2	28.4
ion	ang	26.8	27.6	27.3	27.2	(RCP4.5)	Upper Bound	1.7	28.5	1.8	29.4	1.8	29.1	1.8	29.0
Region 9	mpg	-	2014	50100	31,00	High	Lower Bound	1.4	28.2	1.4	29.0	1.4	28.7	1.3	28.5
1	Za					(RCP8.5)	Median	1.7	28.5	1.8	29.4	1.5	28.8	1.6	28.8
B						(Upper Bound	2.2	29.0	2.3	29.9	2.3	29.6	2.3	29.5
	· P					Moderate	Lower Bound	0.9	28.0	1.1	29.0	0.9	28.4	1.0	28.5
ľ	ang	1	5-3			(RCP4.5)	Median Upper Bound	1.8	28.9	1.7	29.2	1.6	29.1	1.7	29.2
۱	Zamboanga Sibugay	27.1	27.9	27.5	27.5		Lower Bound	1.4	28.5	1.4	29.3	1.3	28.8	1.4	28.9
ľ	Zar					High Emission	Median	1.6	28.7	1.7	29.6	1.4	28.9	1.5	29.0
П	77					(RCP8.5)	Upper Bound	2.2	29.3	2.2	30.1	2.0	29.5	2.2	29.7

^{*}percentile range of model projections: lower bound = 10th percentile, median = 50th percentile, upper bound = 90th percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperature (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

Ī	1.7	Obs	served (1	971-200	00)				Pro	jected (2	036-2065)				
_	9	1	1					DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	МАМ	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
						Moderate	Lower Bound	1.1	26.2	1.1	27.6	1.0	26.8	1.1	26.8
	uc					Emission (RCP4.5)	Median	1.3	26.4	1.3	27.8	1.2	27.0	1.2	26.9
	Bukidnon	25.1	26.5	25.8	25.7	(RCP4.5)	Upper Bound	1.7	26.8	1.8	28.3	1.9	27.7	1.8	27.5
	Bul					High	Lower Bound	1.4	26.5 26.8	1.5	28.0 28.3	1.4	27.2	1.4	27.1
						(RCP8.5)	Median Upper Bound	2.2	27.3	1.8	28.3	2.4	28.2	2.3	28.0
P						CONTRACT OF	Lower Bound	1.0	25.4	1.1	26.6	1.0	26.4	1.0	26.2
	Norte					Moderate Emission	Median	1.3	25.7	1.3	26.8	1.2	26.6	1.2	26.4
	del N	24.4	25.5	25.4	25.2	(RCP4.5)	Upper Bound	1.8	26.2	1.8	27.3	1.8	27.2	1.8	27.0
Н	9 0	24.4	25.5	25.4	25.2	High	Lower Bound	1.5	25.9	1.4	26.9	1.3	26.7	1.3	26.5
10	Lanao					Emission (RCP8.5)	Median	1.7	26.1	1.8	27.3	1.5	26.9	1.6	26.8
Region						(KCP0.5)	Upper Bound	2.2	26.6	2.3	27.8	2.3	27.7	2.3	27.5
Reg	ntal					Moderate	Lower Bound	1.0	26.6	1.0	27.7	0.9	27.5	1.0	27.4
П	Misamis Occidental		100	17.5		(RCP4.5)	Median Upper Bound	1.3	26.9 27.2	1.2	27.9	1.2	27.8 28.4	1.2	27.6 28.2
	ő	25.6	26.7	26.6	26.4		Lower Bound	1.4	27.0	1.4	28.1	1.3	27.9	1.3	27.7
	amis		L 1			High Emission	Median	1.6	27.2	1.7	28.4	1.6	28.2	1.6	28.0
	MIS					(RCP8.5)	Upper Bound	1.9	27.5	2.1	28.8	2.3	28.9	2.3	28.7
P	_					Moderate	Lower Bound	1.0	26.4	1.0	27.8	1.0	27.9	1,0	27.5
	Misamis Oriental					Emission	Median	1.3	26.7	1.2	28.0	1.2	28.1	1.1	27.6
	ő	25.4	26.8	26.9	26.5	(RCP4.5)	Upper Bound	1.7	27.1	1.7	28.5	1.8	28.7	1.8	28.3
	SE	25.4	20.0	20.2	20.5	High	Lower Bound	1.4	26.8	1.4	28.2	1.3	28.2	1.3	27.8
	Misa					(RCP8.5)	Median	1.6	27.0	1.7	28.5	1.6	28.5	1.5	28.0
-	(2)					(ACFO.3)	Upper Bound	2.1	27.5	2.3	29.1	2.2	29.1	2.2	28.7
	lley					Moderate	Lower Bound	1.0	27.7	1.1	28.9	1.0	28.6	1.0	28.6
	Compostella Valley			2000		(RCP4.5)	Median Upper Bound	1.7	28.4	1.8	29.1	1.8	29.4	1.8	29.4
	stell	26.7	27.8	27.6	27.6		Lower Bound	1.4	28.1	1.5	29.3	1.3	28.9	1.3	28.9
	odu		1			High Emission	Median	1.6	28.3	1.7	29.5	1.7	29.3	1.6	29.2
	Ö					(RCP8.5)	Upper Bound	2.2	28.9	2.3	30.1	2.3	29.9	2.2	29.8
						Moderate	Lower Bound	1.0	27.7	1.0	28.8	1.0	28.4	1.0	28.4
	lorte					Emission	Median	1.2	27.9	1,2	29.0	1.2	28.6	1.2	28.6
	Davao del Norte	26.7	27.8	27.4	27.4	(RCP4.5)	Upper Bound	1.8	28.5	1.8	29.6	1.8	29.2	1.8	29.2
	900					High	Lower Bound	1.4	28.1	1.5	29.3	1.3	28.7	1,3	28.7
	Dav					(RCP8.5)	Médian	1.6	28.3	1.7	29.5	1.6	29.0	1.6	29.0
1						(Marais)	Upper Bound	1.0	28.9 27.9	2,3	30.1	2.3	29.7	2.2	29.6
۵	Sur					Moderate Emission	Lower Bound Median	1.0	28.1	1.0	28.8	1.0	28.1	1.0	28.1
11	e s	270.0	100	170.00		(RCP4.5)	Upper Bound	1.7	28.6	1.7	29.5	1.8	28.7	1.8	28.9
Region	90 d	26.9	27.8	26.9	27.1	unac	Lower Bound	1.3	28.2	1,4	29.2	1.3	28.2	1,3	28.4
æ	Davao del					High Emission	Median	1.6	28.5	1.7	29.5	1.6	28.5	1.6	28.7
						(RCP8.5)	Upper Bound	2.2	29.1	2.3	30.1	2.3	29.2	2.2	29.3
	To					Moderate	Lower Bound	1.0	27.8	1.0	28.8	1.0	28.5	1.0	28.6
	Occidental					(RCP4.5)	Median	1.2	28.0	1.2	29.0	1.2	28.7	1.3	28.9
	Occi	26.8	27.8	27.5	27.6	(1007-1.3)	Upper Bound	1.6	28.4	1.7	29.5	1.8	29.3	1.8	29.4
	080		100			High Emission	Lower Bound Median	1.3	28.1	1.3	29.1	1.3	28.8	1.3	28.9
	Dav					(RCP8.5)	Upper Bound	2.0	28.4	2.1	29.4	2.2	29.0	2.2	29.1
0						1200000	Lower Bound	1.0	27.8	1.0	28.8	1.0	28.5	1.0	28.6
	E					Moderate Emission	Median	1.2	28.0	1.2	29.0	1.1	28.6	1.2	28.8
	Davao Orienta	20.0	22.5	22.5	200	(RCP4.5)	Upper Bound	1.7	28.5	1.7	29.5	1.7	29.2	1.8	29.4
	30 0	26.8	27.8	27.5	27.6	High	Lower Bound	1.3	28.1	1.4	29.2	1.3	28.8	1.3	28.9
	Dav					Emission	Median	1.6	28.4	1.6	29.4	1.6	29.1	1.5	29.1
	1	11.				(RCP8.5)	Upper Bound	2.1	28.9	2.3	30.1	2.1	29.6	2.2	29.8
	o					Moderate	Lower Bound	1.0	27.8	1.0	28.9	1.0	28.0	1.0	28.1
	North Cotabato		1 1			(RCP4.5)	Median	1.3	28.1	1.2	29.1	1.2	28.2	1.2	28.3
	Cot	26.8	27.9	27.0	27.1	(10)	Upper Bound	1.8	28.6	1.7	29.6	1.8	28.8	1.8	28.9
	ŧ		1000			High Emission	Lower Bound	1.4	28.2 28.6	1.4	29.3	1.3	28.3 28.6	1.4	28.5 28.7
112						(RCP8.5)	Median Upper Bound	2.3	29.1	2.3	30.2	2.3	29.3	2.3	29.4
Region		11 11				46.6	Lower Bound	1.0	28.7	1.1	29.5	1.0	28.3	1.0	28.6
Re	-					Moderate Emission	Median	1.2	28.9	1.3	29.7	1.2	28.5	1.3	28.9
	gan	22.7	20.4	27.0	27.5	(RCP4.5)	Upper Bound	1.7	29.4	1.7	30.1	1.8	29.1	1.8	29.4
	Sarangani	27.7	28.4	27.3	27.6	High	Lower Bound	1.4	29.1	1.4	29.8	1.4	28.7	1.3	28.9
	N.					Emission	Median	1.6	29.3	1.7	30.1	1.6	28.9	1,6	29.2
						(RCP8,5)	Upper Bound	2.1	29.8	2.3	30.7	2.2	29.5	2.2	29.8

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile



Table B-2. CLIRAM of the projected seasonal change in mean temperature (in degree Celsuis) in the mid-21st century (2036-2065) for all provinces; baseline period: 1971-2000 (con't).

		Obs	served (1	971-20	00)				Pro	jected (2	036-2065)				
_	8			100				DJF (Dec	-Jan-Feb)	MAM (Mai	-Apr-May)	JJA (Jun	-Jul-Aug)	SON (Sep	-Oct-Nov)
Region	Province	DJF	МАМ	JJA	SON	Scenario	Range*	Change	Projected value	Change	Projected value	Change	Projected value	Change	Projected value
	10.					Moderate	Lower Bound	1.0	28.7	1.1	29.6	1.1	28.5	1.0	28.7
	Cotabato					Emission	Median	1.3	29.0	1,3	29.8	1.2	28.6	1.3	29.0
	otat	22.2	20.5	27.4	27.7	(RCP4.5)	Upper Bound	1.8	29.5	1.8	30.3	1.9	29.3	1.9	29.6
	0	27.7	28.5	27.4	21.1	High	Lower Bound	1.4	29.1	1.5	30.0	1.4	28.8	1.4	29.1
7	South					Emission	Median	1.7	29.4	1.8	30.3	1.6	29.0	1.6	29.3
n 1	U1					(RCP8.5)	Upper Bound	2.3	30.0	2.4	30.9	2.4	29.8	2.3	30.0
Region	Hy					Moderate	Lower Bound	1.0	28.8	1.0	29.6	1.0	28.6	1.0	28.8
č	arat					Emission	Median	1.2	29.0	1.2	29.8	1.1	28.7	1.2	29.0
	pny	27.0	20.6	27.6	27.8	(RCP4.5)	Upper Bound	1.8	29.6	1.8	30.4	1.8	29.4	1.8	29.6
	- Le	27.8	28.6	27.6	27.8	High	Lower Bound	1.3	29.1	1.4	30.0	1.3	28.9	1.3	29.1
	Sultan Kudarat					Emission	Median	1.7	29.5	1.8	30.4	1.6	29.2	1.6	29.4
						(RCP8.5)	Upper Bound	2.3	30.1	2.3	30.9	2.3	29.9	2.2	30.0
Ī	aı					Moderate	Lower Bound	1.0	27.2	1.1	28.7	0.9	28.7	1.0	28.4
	Norte					Emission	Median	1.2	27.4	1.3	28.9	1.2	29.0	1.2	28.6
	- E	20.2	27.6	27.0	27.4	(RCP4.5)	Upper Bound	1.7	27.9	1.8	29.4	1.8	29.6	1.8	29.2
	Agusan del	26.2	27.6	27.8	27.4	High	Lower Bound	1.4	27.6	1.5	29.1	1.3	29.1	1.3	28.7
	asnt					Emission	Median	1.6	27.8	1.7	29.3	1.7	29.5	1.6	29.0
	A					(RCP8.5)	Upper Bound	2.1	28.3	2.3	29.9	2.2	30.0	2.3	29.7
	771					Moderate	Lower Bound	1.0	26.9	1.1	28.2	1.0	28.2	1.0	27.9
	Sur					Emission	Median	1.2	27.1	1.2	28.3	1.1	28.3	1.2	28.1
		635	120	202	337	(RCP4.5)	Upper Bound	1.7	27.6	1.8	28.9	1.8	29.0	1.8	28.7
	Agusan del	25.9	27.1	27.2	26.9		Lower Bound	1.3	27.2	1.4	28.5	1.4	28.6	1.4	28.3
	snb					High Emission	Median	1.6	27.5	1.7	28.8	1.7	28.9	1.6	28.5
CARAGA	×					(RCP8.5)	Upper Bound	2.2	28.1	2.3	29.4	2.3	29.5	2.2	29.1
8	ni.					the desired	Lower Bound	1.0	27.3	1.0	28.6	0.9	29.1	1.0	28.7
ū	orte					Moderate Emission	Median	1.2	27.5	1.2	28.8	1.2	29.4	1.1	28.8
	del Norte	123	-33.52	100		(RCP4.5)	Upper Bound	1.7	28.0	1.6	29.2	1.7	29.9	1.8	29.5
	0	26.3	27.6	28.2	27.7		Lower Bound	1.3	27.6	1.4	29.0	1.2	29.4	1.3	29.0
	Surigao		10.00			High Emission	Median	1.5	27.8	1.6	29.2	1.6	29.8	1.5	29.2
	Su					(RCP8.5)	Upper Bound	2.0	28.3	2.1	29.7	2.1	30.3	2.2	29.9
						4 T/3 T/5	Lower Bound	1.0	27.4	1.0	28.4	1.0	28.9	1.0	28.4
	Sur					Moderate Emission	Median	1.2	27.6	1.2	28.6	1.2	29.1	1.2	28.6
	del	ra.	55.7	1000		(RCP4.5)	Upper Bound	1.7	28.1	1.7	29.1	1.8	29.7	1.8	29.2
	90	26.4	27.4	27.9	27.4		Lower Bound	1.3	27.7	1.4	28.8	1.4	29.3	1.4	28.8
	Surigao					High Emission	Median	1.5	27.9	1.5	28.9	1.7	29.6	1.5	28.9
	N.		1			(RCP8.5)	Upper Bound	2.0	28.4	2.1	29.5	2.2	30.1	2.2	29.6
Ť						1000	Lower Bound	1.1	25.4	1.1	26.5	1.0	26.0	1.0	25.9
	Sur					Moderate Emission	Median	1.3	25.6	1.3	26.7	1.2	26.2	1.2	26.1
	E U		1,50	2.5		(RCP4.5)	Upper Bound	1.8	26.1	1.8	27.2	1.8	26.8	1.8	26.7
	o del	24.3	25.4	25.0	24.9		Lower Bound	1.5	25.8	1.4	26.8	1.4	26.4	1.4	26.3
	Lanao					High Emission	Median	1.7	26.0	1.8	27.2	1.6	26.6	1.6	26.5
Σ						(RCP8.5)	Upper Bound	2.3	26.6	2,3	27.7	2.3	27.3	2.2	27.1
ARMM						10.000.00	Lower Bound	1.0	28.6	1.0	29.3	1.0	28.5	1.0	28.6
٩						Moderate Emission	Median	1.0	28.8	1.2	29.5	1.1	28.6	1.2	28.8
	ana		4.0	19-4		(RCP4.5)	Upper Bound	1.8	29.4	1.8	30.1	1.8	29.3	1.8	29.4
	puir	27.6	28.3	27.5	27.6		Lower Bound	1.4	29.4	1.3	29.6	1.3	28.8	1.3	28.9
	Maguindanao		1			High Emission	Median	1.8	29.4	1.8	30.1	1.6	29.1	1.6	29.2
	2					(RCP8.5)		2.3		2.3		2.3	29.1	2.3	29.2
	-					- 10th parce	Upper Bound		29.9		30.6	2.3	29.8	2.3	29,9

^{*}percentile range of model projections: lower bound = 10^{th} percentile, median = 50^{th} percentile, upper bound = 90^{th} percentile

Republic of the Philippines

Department of Science and Technology

Philippine Atmospheric, Geophysical and Astronomical Services Administration

Science Garden Complex, Agham Road, Diliman, Quezon City

ISBN 978-621-95882-0-1